

PAPER • OPEN ACCESS

Electron-impact ionisation of W^{q+} ions with $1 \leq q \leq 19$

To cite this article: D Schury *et al* 2020 *J. Phys.: Conf. Ser.* **1412** 152011

View the [article online](#) for updates and enhancements.

You may also like

- [Absolute rate coefficients for photorecombination of beryllium-like and boron-like silicon ions](#)

D Bernhardt, A Becker, C Brandau et al.

- [Electron-impact ionization of moderately charged xenon ions](#)

M S Pindzola, S D Loch, A Borovik et al.

- [Electron-impact single ionisation of \$W^{q+}\$ ions: experiment and theory for \$11 \leq q \leq 18\$](#)

D Schury, A Borovik, B Ebinger et al.

The advertisement features a yellow header with the PRIME logo and text: "PACIFIC RIM MEETING ON ELECTROCHEMICAL AND SOLID STATE SCIENCE". Below this, it says "HONOLULU, HI October 6-11, 2024". A green section below contains the text: "Joint International Meeting of The Electrochemical Society of Japan (ECSJ) The Korean Electrochemical Society (KECS) The Electrochemical Society (ECS)". To the right, there are two photographs: one showing two men at a booth, and another showing a group of people networking. A teal background section on the right contains the text: "Early Registration Deadline: September 3, 2024" and "MAKE YOUR PLANS NOW!".

Electron-impact ionisation of W^{q+} ions with $1 \leq q \leq 19$

D Schury^{1,2,3*}, A Becker¹, A Borovik Jr^{4†}, B Ebinger^{3,4}, M F Gharaibeh^{1,5,7}, J Hellhund¹, K Huber⁴, J Rausch¹, K Spruck^{1,8}, F Jin^{4,6}, S Schippers⁴ and A Müller¹

¹Institut für Atom- und Molekülphysik, Justus-Liebig-Universität Gießen, 35392 Giessen, Germany

²Institut des Nanosciences de Paris, Sorbonne Université, 75005 Paris, France

³GSI Helmholtzzentrum für Schwerionenforschung GmbH, 64291 Darmstadt, Germany

⁴I. Physikalisches Institut, Justus-Liebig-Universität Gießen, 35392 Giessen, Germany

⁵Department of Physics, Jordan University of Science and Technology, Irbid 22110, Jordan

⁶Department of Physics, National University of Defense Technology, Hunan 410073, Changsha, China

⁷Department of Mathematics, Statistics and Physics, P.O.Box 2713, Qatar University, Doha, Qatar

⁸Institut für Medizinische Physik und Strahlenschutz, THM, 35390 Giessen, Germany

Synopsis Electron-impact ionisation cross sections of tungsten ions in the energy range from the ionisation threshold to 1000 eV were measured. The experimental cross sections are compared with results from sub-configuration averaged distorted wave calculations. Derived plasma rate coefficients are provided for convenient use in plasma modelling.

Tungsten is considered a suitable material for plasma-facing components in fusion reactors due to its good thermo-mechanical properties and its low sputtering yield, low chemical erosion, low tritium retention and high durability against neutron flux [1]. As these components will be subject to high particle fluxes, tungsten impurities in the fusion plasma cannot be avoided [2]. The tungsten ions are expected to absorb collisional energy of free electrons and emit strongly in the X-ray and EUV energy range [3] which leads to energy loss in the optically thin plasma which cannot reabsorb the radiation. Control and understanding of the conditions in the plasma requires extensive modelling and this, in turn, reliable data on atomic processes. This motivated our present study on electron-impact ionisation of tungsten ions which complements related work on photoionisation [4] and recombination [5] of tungsten ions. We measured absolute cross sections for electron-impact single ionisation of tungsten ions of the charge states 1 to 19 employing the animated-beam technique at the Giessen electron-ion crossed-beams experiment [6, 7]. In addition we performed accompanying sub-configuration averaged distorted wave calculations (SCADW) using the FAC code [8] in order to guide the interpretation of the measured cross sections. For easier access in plasma modelling, plasma rate coefficients are derived from the experimental and theoretical cross sections.

*E-mail: Daniel.Schury@physik.uni-giessen.de

†E-mail: Alexander.Borovik@physik.uni-giessen.de




Figure 1. Cross sections for electron-impact single ionisation of W^{12+} ions: Experimental data (solid line, open symbols with quadrature sum of the statistical and systematic uncertainties as error bars) compared to our results of SCADW calculations (dashed line), comprising direct ionisation (DI, dotted line) and excitation autoionisation (EA, dash-dotted line).

References

- [1] Philipp V 2011 *J. Nucl. Mater.* **415** S2
- [2] Dux R *et al* 2011 *Nucl. Fusion* **51** 053002
- [3] Pütterich T *et al* 2010 *Nucl. Fusion* **50** 025012
- [4] Müller A *et al* 2017 *J. Phys. B* **50** 085007
- [5] Krantz C *et al* 2017 *J. Phys. B* **50** 052001
- [6] Rausch J *et al* 2011 *J. Phys. B* **44** 165202
- [7] Borovik A *et al* 2016 *Phys. Rev. A* **93** 012708
- [8] Gu M F 2008 *Can. J. Phys.* **86** 675

