

A LIST OF MARINE ALGAE FROM SEASHORES OF IRAN (HORMOZGAN PROVINCE)

By

J. Sohrabipour and R. Rabii.

Natural Resources and Animal Research Center of Hormozgan. P.O.Box. 79145-1468, Bandar Abbas, Iran.

ABSTRACT

Seashores of Hormozgan province in south of Iran were studied for collection and recognition of algal flora (1991-1996). 150 species of marine algae were recognized, including 36 Chlorophyta (16 new), 33 Phaeophyta (15 new), 75 Rhodophyta (42 new), 3 Xanthophyta (2 new) and 3 Cyanophyta (2 new). In this study number of algal species from Iranian seashores are increased to 201 species.

INTRODUCTION

Marine algae of Persian Gulf for the first time were studied by Endlicher and Diesen (1845). They recorded 8 species of algae based on Kotschy collection from Kharck island in northwest of the Gulf. Borgesen (1939) recorded 102 species of marine algae from Iraian Gulf which 76 species of them were collected from Iranian seashores and others from Arabian regions of the Gulf. Nizamuddin and Gessner (1970) described 68 species of algae from Iranian seashores and Pakistan seashores based on algal collection which had been gathered in «Meteor» Expedition. 37 species of these algae were collected from seashores of Iran and 25 species of them were new records for algal flora of Iran.

From Arabian regions of the Gulf first report of algae were published by Borgesen (1939) which 27 species of the algae had been collected from Arabian regions of the Gulf. Newton (1955a,, 1955b) published the algal flora of Kuwait and Bahrain sea-

shores. In Nizamuddin and Gessner report's (1970) one species was reported from Abudhabi seashores.

Jones (1986) described and illustrated 67 species of algae from seashores of Kuwait and Persian Gulf. Al-Hassan and Jones (1989) recorded 105 species of algae from Kuwait seashores. Basson *et al* (1989) recorded 88 species of algae from Bahrain seashores. Msheghni and Dugham (1987) published benthic marin algae from Qatar seashores. Basson (1992) published a checklist of algae of the Gulf.

Other publication dealing with phytochemical properties or ecological features of the algae of the Gulf including : El-Mohsen and Khoja (1973), Kamel (1981), McCain (1984), McCain *et al* (1984), Basson & Abbas (1992), Abbas *et al* (1992) and Al-Easa *et al* (1995).

Materials and methods :

The present work carried out in seashores of Hormozgan province and its Islands in

the south of iran. Algal specimens were collected form intertidal regions of the seashores (1991-1996) and preserved in 4% Formalin an seawater solution or pressed as herbarium specimens.

All specimens are deposited in natural Resources and Animal Reserach Center of Hormozgan. In laboratory by using of Olympus photostreomicroscope (model SZH) and Olympus photomicroscope (model BH2), morphological and anatomical features of the specimens were studied and photographed. In present paper 150 species of algae are presented form Iranian seashores of the Persian Gulf which 94 of the algae are new records for Iran and based on Basson checklists (1992), 63 species of the algae are new for the Gulf, Systematic account of the recognised algae are presented as follow :

CHLOROPHYTA ULVACEAE

Entromorpha clathrata (Roth) Greville
Abbott & Hollenberg, 1976,P. 73; Al-Hassan & Jones, 1989 p. 292; Basson,, 1979a p.74 &, 1979b p.28' Borgesen, 1939,p.57; Chapman, 1954,p.420; Lawson & John, 1987,p.54; Millar & Kraft, 1994,p.422; Nizamuddin & Gessner, 1970,p.3; Taylor, 1957,p.63; Taupr, 1960,p.59.

Local distrubution : Hormoz Island, Bostanu, Bandar pole and Bandar Jask.

Entrmorpha compressa (Linn.) Nees.

Abbott & Hollenberg1976, p.72; AL-Hassan & Jones, 1989,p.202; Basson, 1992,p.219; Borgesen, 1939,p.57; Chap-

man, 1954,p.406; Millar & Kraft, 1994,p.422; Nizamuddin & Gessner, 1970,p.3; Papenfuss, 1968,p.11; Taylor, 1960,p.60.

Loc. distr.: Wide speread in seashores.

* **Entromorpha flexuosa (Wulf ex Roth) J.Agradh.**

Abbott & Hollenberg, 1976,p.75; Al-Hassan & Jones, 1989,p.292; Basson, 1979a,p.48; Basson, 1989,p.28; Basson, 1992,p.219; Bliding, 1955,p.235-269 (as E. intermedia); Bliding, 1963,pp.73-79; Lawson & John, 1987,p.55; Millar & Kraft, 1994,p.422; N'yeurt, 1996,p.365; Taylor, 1960,p.61;

Loc.distr.: Larak Island, Qeshm Island, Bandar Abbas and Bandar Jask.

***Entromorpha intestinalis (Linn.) Link.**

Abbott & Hollenberg, 1976,p.76; Al-Hassan & Jones, 1989,p.292; Basson, 1989,p.271; Basson, 1992,p.219; Bliding, 1963,pp.140-142; Chapman, 1954,p.404; Nizamuddin & Gessner, 1970,p.3; Papenfuss, 1986,p.11; Lawson & John, 1987,p.55; Steentoft, 1967; Taylor, 1957,p66; Taylor, 1960,p.62.

Loc.distr.: Hormoz Island, Larak Island, Qeshm Island, Hengam Island and Bandar Charak.

* **Entromorpha liza (Linn.) J. Agardh**

Abbott & Hollenberg, 1976,p.76; Bliding, 1963,pp.127; Millar & Kraft, 1994,p.422; Lawson & John, 1987,p.58; Taylor, 1957,p68; Taylor, 1960,p.65.

Loc.distr.: Qeshm Island and Bandar Tiab.

Ulva Lactuca Linn.

Linneaus 1753; Abbott & Hollenberg, 1976,p.83; Al-Hassan & Jones, 1989,p.293; Basson *et al*, 1989,p.28; Basson, 1992,p.219. Bliding, 1968,pp.540-546; Borgesen, 1939,p.57; Chapman, 1954,p.388; Islam, 1976,p.p.11; Lawson & John, 1987,sp.60 Krishnamurthy & Joshi, 196 ,p.126 ; Millar & Kraft, 1994,p.423 ; Richardson, 1975,p.83; Taylor, 1966,p.142 ; Taylor, 1957,p.74 Taylor, 1960,p.65.

Loc.distr.: Qeshm Island, Hormoz Island, larak Island.

Ulva californica Wille

Abbott & Hollenberg, 1976,p.76; Tanner, 1986,pp.510-520.

Loc.distr.: Abumusa Island. Qeshm Island, Bandar Jask.

The species recently was reported (Sohrabipour & Rabii, 1996,p.96).

Ulva fasciata Dlili

Anand, 1940,p.16; Chapman, 1954, p.396' Krishnmurty & Joshi, 1969,p.127; Lawson & John, 1987,p.59; Millar & Kraft, 1994,p.423; Nizamoddin & Gessner, 1970,p.2; Philips, 1988,p.434; Taylor, 1960,p.66; Taylor, 1966,p.142; Sohrabipour & Rabii, 1966.

Loc.distr.: Qeshm Island, Larak Island
The species recently was reported for flora of Iran (Sohrabipour & Rabii, 1996,p.96)

CLADOPHORACEA.E

*** Lola implex (Dillw.) Hamel**

Chapman, 1954,p.463; Islam, 1976,p.13
Loc.distr. : Bandar Jask.

Chaetomorpha aerea (Dillw.) Kutzing.

Al-Hassan & Johnes, 1989,p.293; Basson, 1979a,p.48; Borgesen, 1939,p.63; Basson *et al*, 1989,p.28; Islam, 1976,p.12; Millar & Kraft, 1994,p.425; Papenfuss, 1968,p.13; Taylor, 1957,p.79; Taylor, 1960,p.72 &, 1966,p.143.
Loc.distr. : Qeshm Island.

*** Chaetomorpha brachygona Harvey.**
Islam, 1976,p.12; Lawson & John, 1987,p.65; Rao & Sreeramulu, 1970,p.25; Taylor, 1960,p.70; Taylor, 1966,p.142.
Loc.distr. : Bandar lengeh.

*** Chaetomorpha californica Collins**

Abbot & Hollenberg, 1976,p.101; Chapman, 1954p.463 (as Lola Californica)
Loc.distr. : larak Island.

*** Chaetomorpha garcilis (Kutz.) Kutzing**

Islam, 1976,p.12; Lawson & John, 1987,p.68; Taylor, 1966,p.142;
Loc.distr.: Larak Island

*** Cladophora fassicularis (Mert.) Kutzing**

Chapman, 1954,p.447; Papenfuss, 1968,p.14; Taylor, 1960,p.91 ; Taylor, 1966,p.144; Loc.distr.: Qeshm Island.

*** Caladophora koiei Borgesen**

Basson, 1979a,p.50; Basson, 1989,p.28; Borgesen, 1939,pp.66-68; Nizamuddin & Gessner, 1970,p.4;
Loc.distr.: Qeshm Island.

Borgesen (1939) reported the species from Arabian area of Persian Gulf therefor the species for Iranian seashores is new record.

* **Cladophora Magdalena Harvey**

Anand, 1940,p.29; Taylor, 1957,p.88;
Loc.distr.: Abumousa Island.

* **Cladophora nitellopsis Borgesen**

Al-Hassan & Jones, 1989,p.294; Basson *et al*, 1989,p.28 ; Borgesen, 1939 ,pp:64-66 ; Basson *et al*, 1989 p.29; Nizamuddin & Gessner, 1970,p.4.

Loc.distr.: Larak Island.

* **Cladophora radiosa(Sohr.) Kutzing**

Chapman, 1954,p.446;
Loc.distr.: Larak Island.

* **Cladophora sericioides Borgesen**

Al-Hassan & Jones, 1989,p.294; Basson , 1979a,p.50; Borgesen, 1939,pp.64-66; Basson *et al*, 1989,p.29; Nizamuddin & Gessner, 1970,p.4.

Loc.distr.: Abumousa Island.

SIPHONOCLADACEAE

* **Cladophora membranacea (C. Ag.)**

Borgesen

Anand, 1940,p.47; Lawson & John, 1987,pp.117-113; Sohrabipour & Rabii, 1996, Taylor, 1966,p.148.

Loc.distr.: Hormoz Island, Qeshm Island
The species was recently reported for flora
algal of Iran (Sohrabipour & Rabiei, 1996,p.98)

* **Cladophoropsis Zollingeri (Kutz.)**
Borgesen

Al-Hassan & Jones, 1989,p.294; Basson , 1979a,p.51; Basson *et al*, 1989,p.29; Borgesen, 1939,p.62; Basson, 1992,p.219.

Loc.distr.: Qeshm Island and Hormoz Island.

VALONIACEAE

* **Dictyosphaeria cavernosa (Forssk.)**
Borgesen

Al-Hassan & Jones, 1989,p.294; Basson , 1979a,p.51; Basson *et al*, 1989,p.29; Basson, 1992,p.219. Dawson, 1956,p.90; Islam, 1976,p.16; N'yeurt, 1996,p.371; Millar & Kraft, 1994,p.431; Papenfuss, 1968,p.17; Sohrabipour & Rabii, 1996,p.100; Taylor, 1960,pp.116-117; Taylor, 1964,p.147;

Loc.distr.: Qeshm Island Bandar lengeh
The species was recently reported from
Iran (Sohrabipour & Rabii, 1996,p.100).

DERBESSIACEAE

* **Bryopsis pennata Lamx.var.secund**
(Harv.) Collin & Harvey

Dawson, 1956,p.73; Lawson & John, 1987,pp.92-93; Lawson & Price, 1969,p.285; Steentoft, 1967,p.110; Taylor, 1960,p.132;

Loc.distr.: Farur Island, Qeshm Island, Hormoz Island and Bandar Jask.

* **Bryopsis pennatula J.Agradh**

Abbott & Hollenberg, 1976,p.113; Chapman, 1954,p.484;

Loc.distr.: Qeshm Island.

* **Bryopsis plumosa (Huds.) C.Agradh**

Chapman, 1954,p.476; Lawson & John, 1987,pp.93; Lawson & Price, 1969,p.285; Millar & Kraft, 1994,p.424; N'yeurt, 1996,p.374; Papenfuss, 1968,p.19; Taylor, 1957,p.94; Verheij & Rein, 1993,p.389;
Loc.distr.: Hormoz Island.

CAULERPACEAE

Caulerpa manorensis Nizamuddin

Basson, 1992,p.220; Coppejans & Meinese, 1988,p.186; Coppejans, 1992,p.395; Nizamuddin & Gessner, 1970,p.3; Verhij & Rein, 1993,p.394.

Loc.distr.: Bandar Jask.

Caulerpa racemosa (Forssk.) J.Agardh var.macrophysa (Kutz.) Taylor.

Rao & Sreeramulau, 1970,p.29; Sohrabipour & Rabii, 1996,p.98; Taylor, 1960,pp.155;

Loc.distr.: Larak Island, Qeshm Island and Hormoz Island.

The species was recently reported for flora of Iran (Sohrabipour & Rabii, 1996,p.98).

Caulerpa racemosa (Forssk.) J.Ag. var.peltata (Lamx.) Eubank

N'yeurt, 1960,p.378; Wynn, 1995,p.330;
= Caulerpa peltata lamx. ; Anand, 1940,pp.34-35; Islam, 1976,p.19; Papenfuss, 1968; Millar & Kraft, 1994;p.438; Taylor, 1957,p.49; Taylor, 1960,p.150; Sohrabipour & Rabii, 1996,p.98;
= Caulerpa racemosa ecad. peltata ; Coppejans, 1992,p.410; Coppejalls & Reine, 1992,p.173; Verheij & Reine, 1993,p.124

Loc.distr.: Abumousa Island, Qeshm Island and Hormoz Island.

The varity was recently reported from Iran as Coulerpa peltata lamx but wynne (1995,p.330) and N'yeurt (1996,p.378) believe that C. racemose var.peltata (Lamx.) Eubank is correct.

*** Caulerpa sertularioides (S.G.Gmelin) Howe forma.farlowii (Weber - Van**

Bosse) Borgesen,

Basson *et al*, 1989,pp.404; Basson, 1992,p.220; Coppejans, 1992,p.404; Nizamuddin & Gessner, 1970,p.3; Taylor, 1960,p.144.

The species was reported by Nizamuddin & Gessner (1970) from Oman sea therefore its is a new record for Iranian region of Persian Gulf.

Caulerpa taxifolia (Vahl.) C.Agardh

Anand, 1940,p.36; Islam, 1976,p.22; Coppejans, 1992,p:406 ; Millar & Kraft, 1994,p.438; Lawson & John, 1987,p.90; Lawson & Price, 1969,pp.291-293; Rao & Sreeramu, 1970,p.29; Taylor, 1967,p.52
= Caulerpa mexicana Sonder ex Kuzing ; Basson, 1992,p.220; Taylor, 1960,p.142
= Caulerpa crassifolia (C.Ag.)
J.Ag:Borgesen, 1939,p.47.

Loc.distr.: Qeshm Island Bandar Jask and Hormoz Island.

Borgesen (1939) reported the species as Caulerpa crassifolia form Iranian seasheres but Basson (1992) listed the Borgesen speicemen as synonym for Caulerpa mexicana. Bassed on Coppejans report (1992) Caulerpa Mexicana is a varity of Caulerpa taxifolia.

*** Caluerpa sp.**

Loc.distr.: Qeshm Island and Bandar Jask
The speciements completly different with other species and unknown carfully.

CODIACEAE

*** Codium papilatum Tseng et Gilbert**

Al-Hassan & Jones, 1989,p.295; Basson, 1992,p.220.

Loc.distr.: Qeshm Island and Larak Island.

UDOTEACEAE

***Avarinvillea calithina* Kraft & Olesn-Stojkovich**

Kraft & Olsen-Stojkovich, 1985,pp.339-345; Millar & Kraft, 1994,p.339.

Loc.distr.: Qeshm Island and Farur Island
The species recently report from Iranian seashores (Sohrabipour & Rabii, 1996).

POLYPHSACEAE

***Acetabularia calyculus* Quoy et Gaimard.**

Al-Hassan & Jones, 1989,p.295 Basson, 1979a,p.53; Basson *et al*, 1989,p.29; Basson, 1992,p.220; Islam, 1976,p.24; Millar & Kraft, 1994,p.439; Papenfuss, 1968,p.27; Solms & Solms-laubach, 1985,p.26;

Loc.distr.: Bandar lengeh, Qeshm Island
The species was recently recorded from Iran (Sohrabipour & Rabii, 1996).

***Acetabularia mobii* Solms-laubach**

Anand, 1940,pp.39-41; N'yeurt, 1996,p.397; Papenfuss, 1968,p.28; Solms-Laubach 1895,p.30;

Loc.distr.: Qeshm Island.

The species recently recorded from Persian Gulf (Sohrabipour & Rabii, 1996).

***Polyphysa parvula* (Solms-laubach) Schanetter (Bula & Meyer).**

Millar & Kraft, 1994,p.439; N'yeurt, 1996,p.396; Wynne, 1995,p.333; Solms-Loubach, 1985,p.29 (as *Acetabularia parvula*).

Loc.distr.: Qeshm Island.

The species was recently recorded as *Acetabularia* sp. form Iranian seashores (Sohrabipour & Rabii, 1996).

PHAEOPHYTA

ECTOCARPACEAE

* ***Bachelotia antillarum* (Grunow) Gerloff.**

Lawson & John, 1987,p.103; Millar & Kraft, 1994,p.9; Richardson, 1975,p.97; Rao & Sreeramulu, 1970,p.31; Steentoft, 1967,p.115; Taylor, 1960,p.198 (as *B. fulvescens* (Bornet) Kuckuck).

Loc.distr.: Qeshm Island and Larak Island.

* ***Feldmania irregularis* (kutz.) Hamel**

Abbott & Hollenberg, 1976,p.136; Basson, 1979a,p.53; (as *Ectocarpus irregularis* Kutz) Basson, 1992,p.220; Boudouresque *et al*, 1984,p.50; Islam, 1976,p.30 (as *Giffordia irregularis* Kutz) ; Taylor, 1960,p.207 (as *Giffordia conifera*); Papenfuss, 1968,p.29;

Loc.distr.: Qeshm Island, Hormoz Island and Larak Island.

The species recently recorded from Iranian seashores (Sohrabipour & Rabii, 1996,p.101).

***Hinksia mitchelliae* (Harvey) P.C.Silva**

Basson, 1992,p.220; Millar & Kraft, 1994,p.10; Wynne, 1995,p.323;
= *Giffordia mitchelliae* : Abbott & Hollenberg, 1976,p.143; Al-Hassan & Jones, 1989,p.296; Basson, 1979a,p.53; Steentoft, 1967,p.115;

= *Ectocarpus mitchelliae* : Borgesen, 1939,p.75; Nizamuddin & Gessner, 1970,p.5;

Loc.distr.: Larak Island, Qeshm Island and Hormoz Island.

Boryesen (1939) and Nizamuddin & Gessner (1970) reported the species as *Ectocarpus mitchelliae* Harvey form Iranian seashores.

SPHACELARIACEAE

Sphacelaria nova-hollandiae sonder

Al-Hassan & Jones, 1989,p.296; Taylor, 1960,pp.211-12; Wynne, 1995,p.323
Loc.distr.: Abumousa Island.

The species recently recorded from Iranian seashores (Sohrabipour & Rabii, 1996,p.101).

Sphacelaria rigidula Kutzing

Al-Hassan & Jones, 1989,p.296; Basson, 1992,p.221; Niekerk, 1965,p.450 N'yeurt, 1996,p.398; Lawson & John, 1987,p.117; Rao & Sreeramulu, 1970,p.31;
= *Sphacelaria furcigera* Kutzing : Basson, 1979a,p.53; Basson *et al*, 1989,p.31; Borgesen, 1939,p.81; Chamberlain, 1965,p.202; Setchell & Gardner, 1976,p.397; Von den Hoek & Flinterman, 1968,p.193-241;

Loc.distr.: Bandar lengeh.

The species was previously reported from Iranian seashores by Borgesen (1939) as *Sphacelaria furcigera* Kutz.

Sphacelaria tribuloides meneghini

Al-Hassan & Jones, 1989,p.296; Basson, 1979,p.55; Basson *et al*, 1989,p.31; Basson, 1992,p.220; Richardson, 1975,p.99; Papenfuss, 1968,p.31; Rao & Sreeramulu, 1970,p.31; Lawson & John, 1987,p.117; Abumusa Island, Sohrabi & Rabii 278
The species recently recorded from Iran (Sohrabipour & Rabii, 1996,p.102).

DICTYOTACEAE

* **Dictyota ciliolata Kutzing**

Al-Hassan & Jones, 1989,p.297; Lawson & John, 1987,p.122; Richardson, 1975,p.101; Papenfuss, 1968,p.32; Verheij

& Rein, 1993,p.424; Taylor, 1960,p.223; Taylor, 1966,p.159;
Loc.distr.: Qeshm Island

* **Dictyota dichotoma (Hudson) Lamx.**

Basson, 1979a,p.55; Basson, 1992,p.221; Islam, 1976,p.36; Lawson & John, 1987,p.123; Papenfuss, 1968,p.32; Taylor, 1960,p.218; Verheij & Rein, 1993,p.425; Wynne, 1995,p.326;
Loc.distr.: Qeshm Island and Bandar Charak

* **Dictyota divaricata Lamx.**

Allender & Kraft, 1983,p.112; Al-Hassan & Jones, 1980,p.297; Basson, 1979a,p.55; Basson *et al*, 1989,p.31; Basson, 1992,p.221; Borgesen, 1939,p.81; Islam, 1976,p.36; Lawson & John, 1987,p.124; Millar & Kraft, 1994,p.17; Nizamuddin & Gessner, 1970,p.33; Papenfuss, 1968,p.33; Taylor, 1966,p.158.

Loc.distr.: Qeshm Island

The species had recorded from Iran by Borgesen (1939).

* **Dictyota indica Sonder et Kutzing**

Basson, 1992,p.221; Nizamuddin & Gessner, 1970,p.5; Taylor, 1960,p.221; Taylor, 1966,p.158.

Loc.distr.: Qeshm Island.

* **Dictyota volubilis Kutzing sensu Vickers**

Taylor, 1960,p.220; Taylor, 1966,p.158;
Loc.distr.: Qeshm Island

Lobophora variegata (Lamx.) Womersley.

Allender & Kraft, 1983,pp.81-82; Basson,

1992,p.221; Millar, 1990,p.469; Millar & Kraft, 1994,p.18; Lawson & John, 1987,p.127; Islam, 1976,p.38; Papenfuss, 1963,p.35-36; Richardson, 1975,p.102; N'yeurt, 1996,p.401; Womersley, 1967,p.221; Verheij & Rein, 1993,p.426;
= *Pocockiella variegata* (Lamx.) Papenfuss ; Basson, 1979a,p.57; Rao & Sreeramulu, 1970,p.32; Dickimson & Foote, 1951,p.137; Taylor, 1960,p.231;
= *Zonaria variegata* (Lamx.) C.Agardh ; Borgesen, 1939,pp.82-83;

Loc.distr.: Qeshm Island, Abumusa Island and Bandar lengeh

Borgesen (1939) recorded the species as *Zonaria variegata* form Iran

* ***Padina australis* Hauck**

Al-Hassan & Jones, 1989,p.297; Allender & Kraft, 1983,p.88; Millar, 1990,p.19; Lawson & John, 1987,p.128; Verheij & Rein, 1993,p.428;

= *Padina gymnospora* (Kutz.) Cickers : Basson, 1979,p.55; Basson *et al*, 1989,p.31; Islam, 1976,p.403; Nizamuddin & Gessner, 1970,p.6; Richardson, 1975,p.104; Taylor, 1960,p.236; Taylor, 1966,p.161;

Loc.distr.: Qeshm Island.

* ***Padina borgesenii* Allender & Kraft**

Allender & Kraft, 1983,pp.37-89; Millar & Kraft, 1994,p.19; Verheij & Rein, 1993,p.429;

Loc.distr.: Qeshm Island.

* ***Padina crassa* Yamada**

Allender & Kraft, 1983,p.87; Millar, 1990,p.470;
Millar & Kraft, 1994,p.19;

Loc.distr.: Larak Island.

* ***Padina pavonica* (Linn.) Thivy**

Islam, 1976,pp.40-41; Lawson & John, 1987,p.132; Papenfuss, 1986,p.34; Taylor, 1960,p.234;

= *Fucus pavonica* linn. : Linnaeus 1753,2:1162

= *Padina cf. pavonia* (Linn.) Baill : Kunkel, 1977,p.156;

Loc.distr.: Qeshm Island.

The species hade been uncertainly reported as *P. pavonia* (Linn) Baill. from Hormoz Island by Kunkel (1977).

* ***Padina Tenuis* Bory de saint vincent**

Allender & Kraft, 1983,p.83; Millar & Kraft, 1994,p.19; Verheij & Rein, 1993,p.429; N'yeurt, 1996,p.402;

= *Padina boryana* Thivy : Basson, 1992,p.221;

= *Padina commersonil* Bory : Borgesen, 1939,p.82; Papenfuss, 1968,p.34; Steentoft, 1967,p.118;

Loc.distr.: Bandar Charak, Qeshm Island

Borgesen (1939) recorded the species as *padina commersonil* Bory and Basson (1992) listed as synonym for *Padina boryana* Thivey, but based on Allener & Kraft (1983) both two species (*P. commersonil* & *P. boryana*) are synonym *P. tenuis* Bory de Saint Vincent.

***Padina tetrastromatica* Hauck**

Allender & Kraft, 1983,pp.86-87; Basson, 1992,p.221; Borgesen, 1939,pp.81-82; Dickimson & Foote, 1951,p.134; Islam, 1976,p41; Lawson & John, 1987,p.131; Newton, 1953,p.396; Papenfuss, 1968,p.34; Rao *et al*, 1994,p.43-46;

Loc.distr.: Larak Island and Qeshm Island

The species perviously recorded by Borgesen (1939) form Iran.

* **Spatoglossum variables** Figail et De-notaris

Nizamuddin & Gessner, 1970,p.6; Papenfuss, 1968,p.35;

Loc.distr.: Qeshm Island and Larak Island

Nizamuddin & Gessner (1970) recorded the species from Pakistan seashores in Oman sea therefor it is a new record for Iranian seashore of Persian Gulf.

* **Spatoglossum marginatum** (C. Ag.)
Kutzing

Papenfuss, 1968,p.35; Phillips *et al.*, 1993,pp.395-398;

Loc.distr.: Qeshm Island, Bandar Lengeh, Bandar Charak and bandar Jask.

PUNCTARICACEAE

Colpomenia sinuos (Mertens ex Roth)

Derbes et Solier

Al-Hassan & Jones, 1989,p.297; Basson, 1979a,p.57; Basson, 1992,p.221; Basson *et al.*, 1989,p.38; Borgesen, 1939,p.89; Chamberlain, 1965,p.203; Dawson, 1956,p.91; Dickimso & Foote, 1951,p.137; Lawson & John, 1987,p.137; Millar & Kraft, 1994,p.22; Nizamuddin & Gessner, 1970,p.6; Papenfuss, 1968,p.38; Steentoft, 1967,p.120; Taylor, 1960,p.260 &, 1966,p.162; Vandermeulent *et al.* =, 1984

= Asperococcus sinuosus Bory var.lobutus: Endlicher & Diesing 1845.
Loc.distr.: Qeshm Island, Hengam Island. Larak Island, Bandar Charak and Bandar lengeh.

The species had been reported from Iranian seashores by Endlicher & Diesing (1845) as Asperococcus sinuosus Bory var.lobutus and by Borgesen (1939).

Lyengaria stellata (Borgesen) Borgesen
Al-Hassan & Jones, 1989,p.297; Basson, 1992,p.221; Borgesen, 1939,pp.91-93; Nizamuddin & Gessner, 1970,p.6;
Loc.distr.: Qeshm Island, Larak Island, Hengam Island and Bandar Charak Borgesen (1939) and Nizamuddin & gessner (1970) recorded the species form Iranian seashores.

* **Rosenvingea floridana** (Tylor) Taylor
Abbott & Hollenberg, 1976,p.220; Taylor, 1960,p.262; Taylor, 1964,p.162;
Loc.distr.: Qeshm Island and Bandar Charak.

CYSTOSEIRACEAE

Cystoseira myrica (S. G . Gmlin)

C.Agardh

Al-Hassan & Jones, 1989,p.298; Basson, 1970a,p.57; Basson, 1992,p.221; Basson *et al.*, 1989,p.31; Borgesen, 1939,p.95; Endlicher & Diesing 1845 (as C.myrica.var.tenella Hering et Martens); Newton, 1955a,p.101; Newton, 1955,p.142; Papenfuss, 1968,p.40; Taylor, 1960,p.267; Taylor, 1966,p.163;
Loc.distr.: Bandar lengeh, Qeshm Island, Bandar Charak, Bandar Gavbandy and Larak Island.

Cystoseira trinodis (Forskal) C.Agardh

Al-Hassan & Jones, 1989,p.298; Basson, 1979a,p.57; Basson *et al.*, 1989,p.32; Jones, 1986; Papenfuss, 1968,p.41;
= Cystophyllum muricatum (C. Ag.)
J.Agardh:Baorgesen, 1939,p.95; Nizamuddin & Gessner, 1970,p.6; Newton, 1955,p.101 &, 1955b,p.14

= Cystoseira viragta Endlicher et Diesing ;

Endlicher & Diesing 1845

Loc.distr.: Bandar Charak, Qeshm Island and Bandar Lengeh

The species had been reported with different name form Iran by Endlicher & Diesing 1845 (as Borgesen (1939) & Nizamuddin & Gessner (1970).

*** *Cystoseira neglecta* Setchell & Gardner**

Abbott & Hollenberg, 1976,p.269; Setchell & Gardner, 1967,p.710;
Loc.distr.: Bandar Jask.

*** *Hormophysa cuneiformis* (J.G .Gmelin) P.C.Silva**

Al-Hassan & Jones, 1989,p.298; Basson, 1979a,p.59; Basson *et al*, 1989,p.32; Newton, 1955b,p.142; Papenfuss, 1968,p.41; Verheij & Rein, 1993,p.431

= *Hormophysa triquetra* (C. Ag.) Kutzning : Borgessen, 1939,p.96

Loc.distr.: Bandar lengeh

Boegesen (1939) in his report from Iranian Gulf recorded the species from Arabian regions of the Gulf.

SARGASSACEAE

***Sargassum angustifolium* C. Agardh**

Al-Hassan & Jones, 1989,p.298; Basson, 1979a,p.59; Basson, 1992,p.221; Borgesen, 1939,p.100(as *S. flexile* Greville); Endlicher & Diesing 1845,p.268; Jones, 1986; Newton, 1955a,p.101 &, 1955b,p.142; Papenfus, 1968,p.43;
Loc.distr.: Bandar Charak and Qeshm Island.

Nizamuddin & Gessner (1970) reported the species from Iran.

***Sargassum boveanum* J. Agardh**

Al-Hassan & Jones, 1989,p.97; Basson, 1979a,p.59; Basson *et al*, 1989,p.32; Borgesen, 1939,p.97; Basson, 1992,p.222; Jones, 1986; Newton, 1955b,p.142; Nizamuddin & Gessner, 1970,p.7; Papenfuss, 1968,p.45;

Loc.distr.: Qeshm Island.

*** *Sargassum fluitans* (Borgesen)**

Borgesen

Basson, 1992,p.222; Kamel, 1981; Taylor, 1957,p.198; Taylor, 1960,p.281; Taylor, 1964,p.164;

Loc.distr.: Qeshm Island.

*** *Sargassum latifolium* (Turner) C. Agardh**

Basson, 1979a,p.60; Basson *et al*, 1989,p.32;
Basson, 1992,p.222; Papenfuss, 1968,p.52
Loc.distr.: Qeshm Island.

***Sargassum vulgar* C. Agardh**

Basson, 1992a,p.222; Endlicher & Diesing 1845; Islam, 1976,p.47; Lawson & John, 1987,p.142; Rao & Sreeramulu, 1970, p.33; Richardson, 1975,p.106; Steentoft, 1967,p.120; Taylor, 1960,p.242; Taylor, 1966,p.163;

Loc.distr.: Qeshm Island, Bandar Lengeh
Endlicher & Diesing (1845) recorded the species form Iran (Kharck Island).

***Turbinaria conoides* (J. Ag) Kutz.var.
conoides Taylor**

Basson, 1979a,p.60; Borton 1891,p.512;
Taylor, 1960; Taylor, 1964,p.480; Verheij & Rein, 1993,p.434

Loc.distr.: Qeshm Island and larak Island

The species recently recorded from iranian seashores (Shorabipour & Rabii, 1996,p.102).

RHODOPHYTA GONIOTRICHACEAE

***Chroodactylon ornatum* (C. Ag.) Basson**

Al-Hassan & Jones, 1989,p.299; Basson, 1979b,p.67; Basson, 1992,p.227; Basson *et al*, 1989,p.34; Bodouresque *et al*, 1984,p.43; Lawson & John, 1987,p.147; Mozzam & Shameel, 1985,p.145 = *Asterocystis ornata* (C. Ag.) Hamel : Borgesen, 1939,p.102; Levring, 1951,p.460; Newton, 1955a,p.101; Newton, 1955b,p.142

Loc.distr.: ephyphyt on *Laurencia* spp. & *Hypnea* spp.

The species had been recorded as *Asterocystis ornata* from seashores of Iran by Borgesen (1939).

* ***Stylocladia alisidii* (Zanardini) Drew**
Ballesteros, 1991,p.19; Bodouresque *et al*, 1984,p.43;

Lawson & John, 1987,p.147; Lawson *et al*, 1990,p.167;

= *Goniotrichum alisidii* (Zan.) Howe : Abbott & Hollenberg, 1976,p.231; Mozzam & Shameel, 1985,p.144; Papenfuss, 1968,p.68; Taylor, 1957,p.202;
= *G.elegans* (Chauh) Lejolis : Levring, 1951,p.461

Loc.distr.: Qeshm Island (epiphyt on other algae).

ERYTHROPELTIDACEAE

***Erythrocladia irregularis* Rosenvinge** Basson, 1979b,p.67; Basson *et al*,

1989,p.34; Basson, 1992,p.224; Lawson & John, 1987,p.148; Gabarye *et al*, 1980b,p.154; Mozzam & Shameel, 1985,p.146; Heerebout, 1968,p.141; Rao & Sreeramulu, 1970,p.33

= *Erythrocladia subintegra* Rosen.: Abbott & Hollenberg, 1976,p.248; Anand, 1943,p.5; Bodouresque *et al*, 1984; Islam, 1976,p.49; Taylor, 1960,p.290; Taylor, 1966,p.166

Loc.distr.: Qeshm Island (epiphyt on other macro algae).

Borgesen (1939) recorded the species form Iran as *Erythrocladia subintegra* Rosenvinge.

***Erythrotrichia carnea* (Dillwyn) Agradh**

Abbott & Hollenberg, 1976,p.286; Al-Hassan & Jones, 1989,p.300; Basson, 1979b,p.67; Basson *et al*, 1989,p.35; Basson, 1992,p.224; Ballesteros, 1991,p.18; Borgesen, 1939,p.101; Borgesen, 1934,p.113; Gabary *et al*, 1980b,p.156; Islam, 1976,p.49; Lawson & John, 1987,p.149; Levring, 1951,p.462; Mozzam & Shameel, 1985,p.147; Papenfuss, 1968,p.68; Taylor, 1957,p.202; Taylor, 1960,p.292; Taylor, 1966,p.166; Whyne, 1995,p.262

Loc.distr.: Qeshm Island

The species recorded form Iranian seashores by Borgesen (1939).

ACROCHAETIACEAE

***Audouinella Saviana* (Meneghini) Woelkerling**

Millar, 1990,p.299; Lawson & John, 1987,p.157

= *Acrochaetium savianum* Meneghini : Basson, 1979b,p.69; Basson, 1992,p.224;

Boudouresque *et al*, 1984

Loc.distr.: Qeshm Island.

GALAXAURACEAE

Galaxaura lapidscens (Ellis et Solander)

Lamx.

Levrинг, 1951,p.513; Papenfuss, 1968,p.72; Taylor, 1960,p.337; Taylor, 1966,p.167

= *Galaxaura flagelliformis* (Kjell.) emend.Borgesen: Borgesen, 1939,p.104; Loc.distr.: Qeshm Island.

The species had been recorded as *G. flagelliformis* from Iran by Borgesen (1939).

* **Galaxaura rugosa (Ellis et Solander)**

Lamx.

Lawson & John, 1987,p.166; Levrинг, 1951,p.513; Papenfuss, 1968,p.72; Steentoft, 1967,p.123; Taylor, 1960,p.340; Taylor, 1966,p.167; Verheij & Rein, 1993,p.440; Wynne, 1995,p.264.

Loc.distr.: Larak Island and Qeshm Island.

* **Scinaia tsingalensis Tseng**

Levrинг, 1951,pp.509-510; Huisman, 1986,pp.282-284.

Loc.distr.: Qeshm Island and Bandar Jask.

GELIDIACEAE

Gelidiella acerossa (Forsk.) J. Feldmann et Hamel

Basson, 1992,p.225; Basson *et al*, 1989,p.35; Borgesen, 1939,p.107; Hatta & Rein, 1991,p.351; Lawson & John, 1987,p.179; N'yeurt, 1996,p.408; Papenfuss, 1968,p.73; Price & Scott, 1992,pp.28-29; Steentoft, 1967,p.124; Richardson, 1975,p.112; Taylor, 1960,p.351; Taylor, 1966,p.169; Verheij & Rein, 1993,p.454; Wynne, 1995,p.266;

Loc.distr.: Qeshm Island, Bandar Charak, Bandar Kong. The species recorded from Iran by Borgesen (1939).

* **Gelidium chilensis (Montagne) Stantlices & Montalva**

Santlices & Montalva, 1983,pp.187-190; Loc.distr.: Qeshm Island and Farur Island

* **Gelidium pusillum (Stakhous) Lejolis**

Al-Hassan & Jones, 1989,p.301; Anand, 1943,p.11; Basson, 1992,p.225; Borgesen, 1939,p.105; Boudouresque *et al*, 1984,p.44; Hatta & Rein, 1991,p.364; Islam, 1976,pp.52-53; Lawson & John, 1987,p.175; Lawson *et al*, 1990,p.163; MacKain, 1984; N'yeurt, 1996,p.410; Go-deh *et al*, 1992,p.17; Papenfus, 1963,p.74; Rao & Sreeramulu, 1970,p.36; Rao *et al*, 1994,p.44; Renfrewet *et al*, 1988,p.3311; Richardson, 1975,p.113; Taylor, 1960,p.354; Taylor, 1966,p.169

Loc.distr.: Hormoz Island, Qeshm Island and Larak Island .

borgesen (1939) recorded the species from Iranian seashores .

CONRALINACEAE

* **Amphiroa fragilissima (Linn.) Lamx.**

Anand, 1943,p.19; Basson, 1979b,p.69; Basson, 1992,p.225; Borgesen, 1934,p.7; Heiba *et al*, 1990; Jagtap, 1992,p.57; Islam, 1976,pp.53-54; Papenfuss, 1968,p.76; Rao & Sreeramulu, 1970,p.38; Richardson, 1975,p.117; Rao *et al*, 1994,p.44; Price & Scott, 1992,pp.45-47; Taylor, 1960,p.403; Taylor, 1966,p.172; Wynne, 1995,p.270

Loc.distr.: Qeshm Island and Larak Island

* **Amphiroa** sp.

Loc.distr.: Qeshm Island .

Hydrolithon farinosum (Lamx.) Penrose et Chamberlain

N'yeurt, 1996,p412; Penrose & Chamberlain, 1993,pp.295-303;

= *Fosliella farinosa* (Lamx.) Howe : Basson, 1979b,p.70; Basson, 1992,p.225; Basson *et al*, 1989,p.35; Ballesteros, 1991,p.17; Jones, 1986; Lawson & John, 1987,p.205; MacCain, 1984; Rao & Sreeramulu, 1970,p.38; Taylor, 1957,p.252; Taylor, 1960,p.338

= *Melobsia farinosa* Lamx. : Borgesen, 1939,pp.108-109; Boudouresque, 1984,p.46; Endlicher & Diesing 1945,p.269; Newton, 1955a,, 1955b; Wilks & Woelkerlin, 1991,p.528

Loc.distr.: Farur Island, Qeshm Island and Bandar Jask.

Borgesen (1939) recorded the species as *Melobsia forinosa* and Basson (1992) listed as *Fosliella farinosa* but Cahmberlain (1993) described *Hydrolithon farinsoum* (Lamx) Penrose & Chamberlian as new combination .

* **Jania adhaerens** Lamx.

Abbott & Hollenberg, 1976,p.418; Anand, 1943,p.19; Godeh *et al*, 1992,p.17; Islam, 1976,p.53; Lawson & John, 1987,p.202; N'yeurt, 1996,p.413; Papenfuss, 1968,p.77; Price & Scott, 1992,pp.48-49; Steentoft, 1967,p.127; Taylor, 1960,p.413; Taylor, 1966,p.172; Wynne, 1995,p.270

Loc.distr.: Qeshm Island, Larak Island, Bandar lengeh and Hengam Island

Jania rubens (Linn.) Lamx

Basson, 1979b,p.70; Basson, 1992,p.25;

Basson *et al*, 1989,p.107; Borgesen, 1934,p.7; Borgesen, 1939,p.107; Godeh *et al*, 1992,p.17; Johansen & Womersley, 1994,p.607; Lawson & John, 1987,p.204; N'yeurt, 1996,p.413; Papenfuss, 1968,p.78; Rao & Sreeramulu, 1970,p.38; Rao *et al*, 1994,p.44; Steentoft, 1967,p.124; Taylor, 1960,p.413

Loc.distr.: Qeshm Island, Larak Island and Bandar lengeh .

The species was recorded from Iran by Borgesen (1939).

GERACILARIACEAE

Garcilaria corticata (J. Ag.) J. Agardh
Al-Hassan & Jones, 1989,p302; Anand, 1943,pp.57-58; Basson, 1992,p.225; Borgesen, 1939,p.110; Jagtap, 1992,p.57; Nizamuddin & Gessner, 1970,p.9; Rao & Sreeramulu, 1970,p.38; Rao *et al*, 1994,p.44

Loc.distr.: Qeshm Island, Hormoz Island and Bandar Gavbndy

The species had been recorded by Borussen (1939) from Iran.

Garcilaria canaliculata Sonder

Papenfuss, 1968,p.86; Withell *et al*, 1994,pp.301-302; Wynne, 1995,p.281;
Loc.distr.: Qeshm Island, Abumousa Island and Bandar lengeh.

Garcilaria foliifera (Forsk.) Borgesen
Al-Hassan & Jones, 1989,p.302; Basson, 1992,p.225; Borgesen, 1939,p.109; Dawson, 1956,p.177; Nizamuddin & Gessner, 1970,p.9; Papenfuss, 1968,p.86; Richardson, 1975,p.122; Taylor, 1957,p.273; Taylor, 1960,p.446; Taylor, 1966,p.176
Loc.distr.: Hormoz Island, Abumousa Island, Bandar lengeh and Qeshm Island.

The species had been recorded from Iranian by Borgesen (1939) and Nizamuddin & Gessner (1970).

Garcilaria salicornia (C. Ag.) Dawson
Al-Hassan & Jones, 1989,p302; Basson, 1992,p.225; Jagtap, 1992,p.57; Verheij & Rein, 1993,p.458; Withell *et al*, 1994,pp.297-301; Wynne, 1995,p.281
= *Coralopsis cacalia* J.Ag.: Borgesen, 1939,p.110
= *Gracilaria cacalia* (J.Ag.) Dawson : Papenfuss, 1968,p.85
Loc.distr.: Qeshm Island
The species had been recorded as *Coralopsis cacalia* from Iranian seashores by Borgesen (1939).

* **Gracilaria longissima (S. G. Gmelin) Steentoft *et al*, 1995**
Steenstort *et al*, 1995,p.117
= *Gracilaria verrucosa* (Huds.) Papenfuss : Abbott & Hollenberg, 1976,p.500; Dawson, 1956,p.42; Gargiulo *et al*, 1992,p.71; Godeh *et al*, 1992,p.17; Lawson *et al*, 1990,p.166; Lawson & John, 1987,p.227; Millar, 1990,p.339; Papenfuss, 1968,p.87; Richardson, 1975,p.121; Steentoft, 1967,p.133; Taylor, 1957,p.273; Taylor, 1960,p.441; Taylor, 1966,p.173; Verheij & Rein, 1993,p.459 .
Loc.distr.: Bandar Gavbandy .

SOLIERIACEAE

Sarconema filiforme (Sonder) Kylin
Anand, 1943,p.52; Basson *et al*, 1989,p.35; Borgesen, 1934,p.11; Millar, 1990,p.346; Papenfuss, 1968,p.87
= *Sarconema furcellatum* Zanardini : Anand, 1943,p.49; Borgesen, 1932,p.126;

Borgesen, 1934,p.10; Borgesen, 1939,p.111; Islam, 1976,p.55; Papenfuss, 1968,p.83
= *Sarconema furcatum* Borgesen : Anand, 1943,p.52; Borgesen, 1934,p.13
Loc.distr.: Qeshm Island, Larak Island and Bandar lengeh
Borgesen (1939) recorded the species as *S. furcellatum* form Iran but based on Millar report (1990) *S. Furcellatum*, *S. Furcatum* and *S. Montageni* are synonymized *S. Filiform*.

* **Solieria anastomosa Gabrielson & Kraft.**
Gabrielson & Kraft, 1984,pp.222-225;
Loc.distr.: Qeshm Island.

* **Solieria filifomis (Kutz.) Gabrielson.**
Gabrielson, 1985,p.278; Lawson & John, 1987,p.235 .
= *Neogardhiella baileyi* (Kutz.) Wynne et Taylor : Abbott & Hollenberg, 1976,p.483
Loc.distr.: Qeshm Island and Bandar Abbas.

* **Solieria roubsta (Greville) Kylin**
Al-Hassan & Jones, 1989,p302; Borgesen, 1934,p.10; Gabrielson & Kraft, 1984,pp.219-222; Millar, 1990,p.345; Srivivasan, 1969,p.15 .
Loc.distr.: Qeshm Island .

GIGARTINACEAE

* **Gigartina acicularis (Roth) Lamx.**
Boudouresque *et al*, 1984,p.44; Lawson & John, 1987,p.233; Rao & Sreeramulu, 1970,p.39; Millar, 1990,p.358; Taylor, 1960,p.473 .
Loc.distr.: Qeshm Island .

FURCELLARIACEAE

- * ***Furcelaria fastigiata* (Hods.) Lamx.**
Anand, 1943,p.44; Taylor, 1957,p.271
Loc.distr.: Qeshm Island.

HYPNIACEAE

- * ***Hypnea cervicornis* J.Agardh**
Dawson, 1956,p.62; Lawson & John, 1987,p.237; Price & Scott, 1992,pp.35-38;
Jagtap, 1992,p.57; Richardson, 1975,p.124; Steentoft, 1967,p.133; Taylor, 1960,p.466;
Taylor, 1966,p.178
Loc.distr.: Abumousa Island and Qeshm Island.

- * ***Hypnea cornuta* (Kutz.) J.Agardh**
Al-Hassan & Jones, 1989,p.302; Basson, 1979b,p.70; Basson *et al*, 1989,p.35; Dawson, 1956,p.62; Nizamuddin & Gessner, 1970,p.10; Papenfuss, 1968,p.88; Taylor, 1960,p.467; Wynne, 1995,p.274
Loc.distr.: Bandar Abbas, Bandar Charak and Qeshm Island.

- * ***Hypnea flageliformis* Greville ex J.Agardh**
Lawson & John, 1987,p.238
Loc.distr.: Bandar Abbas

- * ***Hypnea musciformis* (Wulfen) Lamx.**
Anand, 1943,p.59; Borgesen, 1939,p.112; Borgesen, 1934,p.17; Dawson, 1956,p.61; Godeh *et al*, 1992,p.17; Islam, 1976,p.55; Lawson & John, 1987,p.240; Lawson *et al*, 1990,p.163; Nizamuddin & Gessner, 1970,p.9; Papenfuss, 1968,p.89; Rao & Sreeramulu, 1970,p.39; Richardson, 1975,p.124; Verheij & Rein, 1993,p.462; Steentoft, 1968,p.133; Taylor, 1957,p.271; Taylor, 1960,p.467; Taylor, 1964,p.178
Loc.distr.: Hormoz Island and Qeshm Island

The species had been recorded by Borgesen (1939) from Iran.

* ***Hypnea pannosa* J.Agardh**

- Borgesn, 1939,p.112; Islam, 1976,p.55; Price & Scott, 1992,p.38-40; Verheij & Rein, 1993,p.462; Wynne, 1995,p.274
Loc.distr.: Hengam Island
Borgesn (1939) recorded the species form Iran.

* ***Hypnea spinella* (C.Ag.) Kutz.**

- Lawson & John, 1987,p.242; Millar, 1990,p.352; Papenfuss, 1968,p.89; Price & Scott, 1992,pp.40-45; Richardson, 1975,p.124; Verheij & Rein, 1993,p.462; Steentoft, 1967,p.134; Taylor, 1960,p.465; Taylor, 1966,p.178; Wynne, 1995,p.276
Loc.distr.: Qeshm Island

* ***Hypnea valentia* (Turner) Montagne**

- Anand, 1943,p.60; Basson, 1979b,p.70; Basson *et al*, 1989,p.35; Borgesen, 1939,p.112; Borgesen, 1934,p.17; Borgesen, 1939,p.113; Lawson & John, 1987,p.242; Papenfuss, 1968,p.89; Rao *et al*, 1994,p.44; Wynne, 1995,p.277
Loc.distr.: Larak Island and Qeshm Island
The species had been recorded from Iran by Borgesen (1939).

RHODYMENIACEAE

- * ***Botryocladia liptopoda* (J.Ag) Kylin**
Anand, 1943,pp.60-61; Borgesen, 1939,p.113; Brodie & Guiry, 1988,p.127; Millar, 1990,p.363; Papenfuss, 1968,p.90
Loc.distr.: Qeshm Island
Borgesn (1939) recorded the species from Pakistan seashores and for Iranian seashores of Persian Gulf is new record.

Rhodymenia sp.

Loc.distr.: Qeshm Island and Larak Island

LOMENTARIACEAE

Lomntaria corallicola Borgesen

Borgesen, 1939,pp.113-117; Millar, 1990,p.373; Price & Scott, 1992,pp.63-66; Wynne, 1995,p.285

Loc.distr.: Bandar Abbas .

The species had been recorded as new species from Iran by Borgesen (1939).

* **Lomntaria baileyana (Hrvey) Forlow**

Dawson, 1956,p.99; Taylor, 1960,p.487; Taylor, 1957,p.287; Taylor, 1966,p.179

Loc.distr.: Farur Island

* **Lomntaria** sp.

Loc.distr.: Qeshm Island

Cartilaginous and ragile structure of samples resemble to *L. baileyana* but arched branches and connected point between branches not observed .

CHAMPIACEAE

* **Champia compressa Harvey var. scindica (Harvey) Borgesen .**

Anand, 1943,p.66; Borgesen, 1931,p.117; Millar, 1990,p.371

Loc.distr.: Bandar Jask .

* **Champia globulifera Borgesen**

Anand, 1943,p.69; Basson *et al*, 1989,p.36; Wynne, 1995,p.285

Loc.distr.: Qeshm Island

* **Champia indica Borgesen**

Al-Hassan & Jones, 1976,p302; Wynne, 1995,p.285

Loc.distr.: Qeshm Island

* **Champia parvula (C. Agardh) Harvey**

Abbott & Hollenberg, 1976,p.565; Al-Hassan & Jones, 1989,p303; Anand, 1943,p.67; Borgesen, 1939,p.116; Boudouresque *et al*, 1984,p.45; Borgesen, 1933,p.122; Dawson, 1956,p.94; Islam, 1976,p.57; Lawson & John, 1987,p.251; Millar, 1990,p.371; N'yeurt, 1996,p.417; Nizamuddin & Gessner, 1970,p.10; Newton, 1955a,p.101; Price & Scott, 1992,p.55; Steentoft, 1967,p.134; Taylor, 1957,p.289; Taylor, 1960,p.490; Taylor, 1966,p.17; Verheij & Rein, 1993,p.467; Wynne, 1995,p.283

Loc.distr.: Qeshm Island

Borgesen (1939) recorded the species from Iran.

DELESSERIACEAE

* **Platysiphonia miniata (J. Ag.) Borgesen**

Lawson & John, 1987,p.294; Lawson *et al*, 1990,p.163; Millar, 1990,p.429

Loc.distr.: Larak Island

* **Taenioma nanum (Kutz.) Papenfuss**

Lawson & John, 1987,p.296; Price & Scott, 1992,pp.146; Taylor, 1966,p.182; Wynne, 1995,p.301

Loc.distr.: Qeshm Island

DASYACEAE

* **Dasya baillovia (S. G. Gemelin) Montagne**

Basson, 1992,p.226; Verheij & Rein, 1993,p.442

= *Dasya pedicellata* C.Ag. : Basson, 1979b,p.75; Basson *et al*, 1989,p.38; Boudouresque *et al*, 1984,p.48; Dawson, 1956,p.50; Islam, 1976,p.62; Taylor,

1967,p.326; Taylor, 1960,p.562; Taylor, 1964,p.182

Loc.distr.: Qeshm Island and Larak Island

* **Pogondophorella sp.**

Abbott & Hollenberg, 1976,p.678

Loc.distr.: Qeshm Island and Larak Island

CERAMIACEAE

Anotrichium tenue (C. Ag) Nageli

Basson, 1992,p.226; Basson *et al*, 1989,p.36; Millar, 1990,p.407; Price & Scott, 1992,p.67

= *Griffithsia tenuis* (Harvey) C. Ag. : Al-Hassan & Jones, 1989,p.303; Borgesen, 1939,p.117; Borgesen, 1931,p.11; Papenfuss, 1968,p.94; Taylor, 1957,p.304; Taylor, 1966,p.180

Loc.distr.: Larak Island (epiphyt on *Galaxaura* sp.)

The species had been recorded as *Griffithsia tenuis* C. Ag. from Iran by Borgesen (1939).

Calithamnion cordatum Borgesen

Nizamuddin & Gessner, 1970,p.11; Price & Scott, 1992,p.75; Taylor, 1960,p.507

= *Aglaothamnion cordatum* (Borg.) Feldman-Mazoyer : Abbott & Hollenberg, 1976,p.616; Basson, 1992,p.226; Rao & Sreeramulu, 1970,p.41; Taylor, 1964,p.180

Loc.distr.: Hormoz Island

Nizamuddin & Gessner (1970) reported the species from Iran.

* **Centroceras apiculatum Yamada**

N'yeurt, 1996,p.420; Price & Scott, 1992,pp.79-81

Loc.distr.: Bandar Lengeh, Hormoz Island and Bandar Jask

* **Centroceras bellum Stechel & Gardner**

Lawson & John, 1987,p.267

Loc.distr.: Hormoz Island and Abumousa Island

* **Centroceras clavulatum (C. Ag.) Montagne**

Abbott & Hollenberg, 1976,p.604; Anand, 1943,p.25; Al-Hassan & Jones, 1989,p.303; Basson *et al*, 1989,p.36; Basson, 1979b,p.72; Borgesen, 1939,p.118; Basson, 1992,p.226; Borgesen, 1934,p.18; Chamberlain, 1965,p.227; Dawson, 1956,p.74; Godeh *et al*, 1992,p.16; Islam, 1976,p.59; N'yeurt, 1996,p.421; Millar, 1990,p.390; Nizamuddin & Gessner, 1970,p.11; Papenfuss, 1968,p.92; Rao & Sreeramulu, 1970,p.42; Rao *et al*, 1994,p.44; Richardson, 1957,p.127; Steentoft, 1967,p.135; Taylor, 1960,p.537; Taylor, 1964,p.181

Loc.distr.: Qeshm Island, Hormoz Island and Bandar Jask

The species had been recorded from Iran by Borgesen (1939).

Ceramium fastigiatum (Wulfen ex Roth) Hrvey.

Basson, 1979b,p.72; Basson *et al*, 1989,p.36; Basson, 1992,p.226; Boudouresque *et al*, 1984,p.48; Islam, 1976,p.59; Lawson & John, 1987,p.274; Nizamuddin & Gessner, 1970,p.10; Taylor, 1957,p.309; Taylor, 1960,p.526; Taylor, 1966,p.180

Loc.distr.: Qeshm Island

Nizamuddin & Gessner (, 1970) recorded the species form Iran.

Ceramium flaccidum (Harvey ex Kutz ing) Adrisson

Lawson & John, 1990,p.165; Lawson & John, 1987,p.270; Millar, 1990,p.395; Price & Scott, 1992,pp.89; Wynne, 1995,p.292

= *Ceramium transversale* Collins et Harvey : Basson, 1979b,p.75; Basson *et al*, 1989,p.36; Nizamuddin & Gessner, 1970,p.10

= *C. masonii* Dawson ; Al-Hassan & Jones, 1970,p.303

Loc.distr.: Qeshm Island, Hormoz Island and Bandar Abbas

The species had been previously reported as *C. transversal* from Iran by Nizamuddin & Gessner (1970)

*** Ceramium manrensis Anand**

Anand, 1943,p.28

Loc.distr.: Bandar Jask.

*** Crounia attenuata (C. Ag.) J. Agradh**

Basson *et al*, 1989,p.36; Basson, 1992,p.226; Boudouresque *et al*, 1984,p.48; Papenfuss, 1968,p.94; Price & Scott, 1992,p.117; Taylor, 1966,p.179; Wollaston, 1984,p.295

Loc.distr.: Qeshm Island.

Spyridia filamentosa (Wulfen) Harvey

Abbott & Hollenberg, 1976,p.608; Al-Hassan & Jones, 1989,p.303; Basson, 1979b,p.75; Basson *et al*, 1989,p.36; Basson, 1992,p.226; Boudouresque *et al*, 1984,p.50; Godeh *et al*, 1992,p.19; Lawson & John, 1987,p.282; Papenfuss, 1968,p.95; Price & Scott, 1992,pp.131; Taylor, 1957,p.317; Taylor, 1960,p.539; Taylor, 1966,p.181; Verheij & Rein, 1993,p.442; Richardson,

1975,p.128; Wynne, 1995,p.301
Loc.distr.: Qeshm Island, Bandar Lengeh, Larak Island
Borgesen (1939) recorded the species from Iranian seashores.

RHODOMELACEAE

*** Acanthophora muscoides (Linn.) Bory de Saint-Vincent**

Al-Hassan & Jones, 1989,p.304; Dawson, 1956,p.62; Lawson & John, 1987,p.297; Richardson, 1975,p.134; Steentoft, 1967,p.136; Taylor, 1964,p.186; Verheij & Rein, 1993,p.443

Loc.distr.: Hengam Island, Qeshm Island and Hormoz Island

Acanthophora spicifera (Vahl) Borgesen

Anand, 1943,p.37; Al-Hassan & Jones, 1989,p.304; Basson, 1979b,p.78; Basson *et al*, 1989,p.38; Abbas *et al*, 1992,p.63; Basson & Abbas, 1992,p.60; Nizamuddin & Gessner, 1970,p.13; Rao & Sreeramulu, 1970,p.43; Rao *et al*, 1992,p.57; Richardson, 1975,p.134; Taylor, 1964,p.186; Verheij & Rein, 1993,p.444; Wynne, 1995,p.301

Loc.distr.: Qeshm Island, Bandar Charak, Hengam Island and Larak Island
Nizamuddin & Gessner (1970) recorded the species form Iranian seashores

*** Chondria bernardii Dangeard**

Lawson & John, 1987,p.307

Loc.distr.: Qeshm Island and Larak Island

*** Chondria cornuta Borgesen**

Anand, 1943,p.36; Basson *et al*, 1989,p.38; Borgesen, 1932,p.130; Rao & Sreeramulu, 1970,p.43

Loc.distr.: Qeshm Island and Hormoz Island

Chondria dasypylla (Woodward) C. Agardh

Abbott & Hollenberg, 1976,p.726; Anand, 1943,p.35; Al-Hassan & Jones, 1989,p.305; Basson, 1979b,p.78; Basson *et al*, 1989,p.38; Abbas *et al*, 1992,p.63; Basson & Abbas, 1992,p.60; Borgesen, 1992,p.121; Borgesen, 1934,p.133; Borgesen, 1932,p.132; Bodouresque *et al*, 1984,p.48; Newton, 1955a,p.101; Newton, 1955b,p.143; N'yeurt, 1996,p.430; Taylor, 1957,p.329; Verheij & Rein, 1993,p.444

Loc.distr.: Qeshm Island and Hormoz Island Borgesen (1939) recorded the species from Iranian seashores .

*** Chondria nidifica Harvey**

Abbott & Hollenberg, 1976,p.726

Loc.distr.: Bandar Jask

*** Chondria oppositoclada Dawson**

Abbott & Hollenberg, 1976,p.727; Godeh *et al*, 1992,p.16

Loc.distr.: Qeshm Island

Digenia simplex (Wulfen) C. Agradh

Al-Hassan & Jones, 1989,p.305; Borgesen, 1939,p.121; Dawson, 1956,p.54; Godeh *et al*, 1992,p.17; Lawson & John, 1987,p.310; Newton, 1955b,p.143; Papenfuss, 1968,p.98; Rao *et al*, 1992,p.57; Taylor, 1966,p.184 &, 1955b,p.143

Loc.distr.: Qeshm Island and Bandar Lengeh

The species had been recorded from Iranian seashores by Borgesen (1939).

Horpsiphonia secunda (C. Ag.) Falkenberg

Abbott & Hollenberg, 1976,p.720; Basson *et al*, 1989,p.38; Borgesen, 1939,p.131 (as Herpsiphonia tenella (C. Ag.) Nagl); N'yeurt, 1996,p.431; Lawson & John, 1987,p.314; Millar, 1990,p.451; Price & Scott, 1992,p.175; Rao & Sreeramulu, 1970,p.43; Verheij & Rein, 1993,p.445; Wynne, 1995,p.307

Loc.distr.: Hormoz Island and Bandar Kong

The species had been recorded as H. tenella from Iran seashores by Borgesen (1939).

Laurencia papillosa (C. Ag.) Greville

Al-Hassan & Jones, 1989,p.305; Basson *et al*, 1989,p.38; Abbas *et al*, 1992,p.63; Basson & Abbas, 1992,p.60; Borgesen, 1939,p.118; Bodouresque *et al*, 1984,p.49; Godeh *et al*, 1992,p.18; Lawson & John, 1987,p.320; Papenfuss, 1968,p.99; Price & Scott, 1992,p.188; Rao *et al*, 1992,p.57; Richardson, 1975,p.135; Taylor, 1964,p.186; Verheij & Rein, 1993,p.447; Wynne, 1995,p.312

Loc.distr.: Bandar Lengeh, Bandar Abbas and Bandar Jask.

*** Laurencia pedicularioides Borgesen**

Borgesen, 1933,p.136; Srinivasan, 1969

Loc.distr.: Bandar Jask and Qeshm Island

*** Laurencia intricata Lamx.**

Godeh *et al*, 1992,p.18; Lawson & John, 1987,p.318; Taylor, 1966,p.178

Loc.distr.: Qeshm Island.

*** Laurencia snyderiae Dawson**

Abbott & Hollenberg, 1976,p.738

Loc.distr.: Qeshm Island, Bandar Abbas, Larak Island and Bandar Lengeh and Qeshm Island.

Leveillea jungermannioides (Hering et Martens) Harvey

Basson, 1979b,p.79; Basson *et al*, 1989,p.38; Borgesen, 1939,p.132; Papenfuss, 1968,p.100; Price & Scott, 1992,p.196; Verheij & Rein, 1993,p.448; Wynne, 1995,p.315

Loc.distr.: Qeshm Island, Hormoz Island, Bandar Lengeh and Bandar Charak. The species had been recorded from Iran by Borgesen(1939).

**Polysiphonia scopulorum Harvey
var.villum (C. Ag.) Hollenberg**

Abbott & Hollenberg, 1976,p.692; Basson *et al*, 1989,p.39; Basson, 1979b (as loposiphonia villum); Boudouresque *et al*, 1984 (as P. scopulorum); Chamberlain, 1951,p.231 (as loposiphnia scopulorum (Harv.) Womersley); N'yeurt, 1996,p.433; Millar, 1990,p.445; Price & Scott, 1992,p.211; Womersley, 1979,p.467
Loc.distr.: Hormoz Island .

*** Polysiphonia variegata (C. Ag.) Zannardini**

Anand, 1943,p.39; Al-Hassan & Jones, 1989,p.305; Basson, 1979b,p.79; Basson *et al*, 1989,p.39; Borgesen, 1934,p.26; Nizamuddin & Gessner, 1970,p.13; Zahid *et al*, 1981,pp.198-200 .

Loc.distr.: Qeshm Island and Hegam Island

Nizamuddin & Gessner (1970) recorded the species form Iran.

Tolypiocladia glomerulata (C. Ag.) Schmitz

Islam, 1976,p.65; Price & Scott, 1992,pp.219; Rao *et al*, 1992,p.57; Verheij & Rein, 1993,p.449; Wynne, 1995,p.321 = Rochera glomerulata : Borgesen, 1931,p.17; Nizamuddin & Gessner, 1970,p.13

Loc.distr.: Qeshm Island, Larak Island and Abumousa Island

Nizamuddin & Gessner (1970) recorded the species as Roschera glomeralata.

**XANTHOPHYTA
NITZSHIACEAE**

Nitzschia martiana (C. Ag.) Van Huerck

Basson & Mohamed, 1985,pp.253-257
Ephyt on other algae
The species recently recorded from Iranian seashores of the Gulf. (Sohrabipour & Rabii, 1996,p.102)

THALASSIOSIRACEAE

*** Stephanopyxis palmeriana (Grev.) Grnow**

Bold *et al*, 1987,p.149

TABELARIACEAE

*** Tabelaria flocculosa (Roth) Kutzing**

Koppen, 1975,pp.236-244

Ephyt on other algae.

**CYANOPHYCEAE
NOSTOCACEAE**

*** Calothrix scopulorum (Weber et Mohr) C. Ag. ex Borner et Flahant**

Islam, 1976,p.74

OSILATORIACEAE

* **Microcoleus lyngbyaceus Kutzing**

Lawson & John, 1987,p.341

Spirolina subsalsa Orested ex Gomount

Moghadam, 1974,p.81; Anand, 1937,p.163; Lawson & John, 1987,p.348

DISCUSSION :

The present work recorded 150 species of marine algae form Hormozgan province seashores in south of Iran. 94 species the algae are new records for Iran and 63 are new records for the Persian Gulf. The species are Chlorophyta 36 species (27 new for Iran and 20 new for Gulf), Phaeophyta 33 species (18 new for Iran and 8 new for Gulf), Rhodophyta 75 species (44 new for Iran and 32 new for Gulf), Xanthophyta 3 species (all new for Iran and 2 new for Gulf), and Cyanophyt 3 species (2 new for Iran and 1 new for Gulf), Camparision of the results wiht Basson checklist (1992) show 63 species are new Gulf, therefore 207 species in Basson checklist increasing to 270 species (Table 1). Camparing of the results with other authors is shown in table 2 which show about two time increasing to Borgesn report from Iranian seashores (1939). 16 species of the new the algae were preciously published (Sohrabipour & Rabii, 1996).

REFERANCE

- 1) **Abbas, J.A. P.W. Basson & A.Y. Gamal El-Din 1992.** Protein contont of benthic marine algae form Bahrain coastline, Indian J. Mrine sciences vol.21:62-63 .
- 2) **Abbott, I.A. & G. J. Hollenberg 1976.** Marine algae of California, Stanford California .
- 3) **Abbott,I.A & R.E. Norris 1993.** new species of Ceramiaceae (Rhodophyta) from the Hawaiian Island. Phycologia. vo. 32(6): 451-461 .
- 4) **Al-Easa,H.S., J.M. Kornprobst & A.M. Rizk 1995** Major sterol composition of samealgae from Qatar : Phytochemistry, vol 39(2) ; 373-374 .
- 5) **Al-Hassan, R.H. & W.E. Jones 1989,** Marrine algal flora nd segreasses of the coast of Kuwait. J.Univ. Kuwai. (Sci) vol.16:289-339 .
- 6) **Allender, B.M. & C. T. Kraft 1983.** The Marine algae of Lord Howe islnad (New south wales) : The dictyotales and Cutleriales (Phaeophyta). Brunonia vol. 6: 73-130 .
- 7) **Anand, P.L. 1940** Marine aglae form karachi, part 1. Chloropyceae, Panij. Univ. Bot. Pub.52 pp.
- 8) **Anand, P. L. 1943.** Marine algae form karachi, part 2. Rhodophyceae.lbid. 73pp.
- 9) **Ananad, P.L. 1973.** An Ecological study of the algae of the British Chalk-cliffs, Part1, J.Ecology, vol.25:153-187 .
- 10) **Athanasiadis, A 1996** Morphology and classification of the Ceramioideae (Rhodophyton) based on phylogentic priciples. Opera Bot.No.128;1-217 .
- 11) **Ballesteros, El. 1991.** Structure of a deep-water community of halimedatuna (Chlorophyceae, Caulerpales) from the North western. mediterranean Collectanea. Bot.(Barcelona) vol.20:5-21

- 12) Basson, P.W. 1972a. marine Algae of the Arabian Gulf coast of Saudis Arabia (First half) Bot.mar.Volxx11:44-46.
- 13) Basson,P.W., 1979b. marine Algae of the Arabian Gulf coast of Saudi Arabia (second half) Bot.Mar.Vol.22:65-82.
- 14) Basson. O. W. & A. Abbas 1992. Elemental composition of some Marine algae from the Bahrain coastlline (Arabian Gulf). Ind.J.Mar.Sci.Vo.21:56-61.
- 15) Basson. p. W & S. A. Mohamed & D. K. Arora, 1989. A survey of the Benthic Marine algae of Bahrain. Bot.Mar.vol.32:27-40 .
- 16) Basson P.W. & S.A.Mohamed 1939 The Tube-dwelling Ditom,Nitzschia martiana(C.Ag.) Van Hurck as recorded from Bahrain (Arabian Gulf). Bot.Mar. Vol.32:253-257 .
- 17) Barton, E.S. 1891. A systematic and structural Account of the genus Turbinari Lamx. Tarns.Linn. Sci. Secon series.vol.11:215-228 .
- 18) Bliding, C. 1955 Entromorpha intermedia, new species of the coasts of sweden, England and wales, Botniska Notiser Vol.108:253-262 .
- 19) Bliding, C. 1960 A preliminary report on some new Mediterranean Green algae Botaiska Notise vol. 113:172-184
- 20) Bliding, C. 1963 A critical survey of European Taxa in Ulaves : Part 1:Compsosiphon, Percursaria, Blidin-gea, Entromorpha. Opera bot. vol.8 (3).
- 21) Bliding, C. 1968. A critical survey of European Taxa in Ulvales 11, Ulva, Ulvaria, Monostroma, Kornmannio lbd. 535-629 .
- 22) Bold, H. C, C. J. Alexopoulos & T. Deleroryas 1987. Morphology of plants and fungi .
- 23) Brodie, J. & M. D. Guiry 1988. Life History and reproduction of Botryo-cladia ardreana sp.nov. (Rhodymeniales, Rhodophyta) from portugal.Phycologia vol.27:109-130 .
- 24) Borgesen, F. 1939. Marine algae from the Iranian Gulf,specially from innermost part near Bushire and the Island Kharg. Danish. Sci.investigation in Iran.part 1.
- 25) Borgesen, F. 1934. Some Indian Rhodophyceae espacially from the shores of the presidency of Bombay IV. Kew. Bull No.1:1-31 .
- 26) Borgesen, F. 1933. Some Indian Rhodophyceae espacially from the shores of the presidency ofBombay III. Kew. Bull No.113-142 .
- 27) Borgesen, F. 1933. Some Indian Rhodophyceae espacially from the shores of the presidency of Bombay IV. Ibid NO.2:1-34 .
- 28) Borgesen, F. 1932. Some Indian Rhodophyceae espacially from the shores of the presidency of Bombay ll. Kew. Bull No,3:113-134 .
- 29) Borgesen, F. 1931. Some Indian Rhodophyceae espacially from the shores of the presidency of Bombay Kew. Bull No.1:1-24 .
- 30) Boudouresque, C.F. , M.P. Boudouresqe & M. Knoepffler-peguy .1984. Inventair des Algues marines Benthoues dansles Pyrenees - orientales (Mediterane - France) vie et Milieu 34 (1):41-59 .

- 31) **Chapman, V. J. 1954.** The marine algae of Newzealand part 1. Myxophyceae and Chlorophyceae. J. Linn. Soci Bot.vol.IV: 323-501 .
- 32) **Chapman, V. J. 1955.** The marine algae of Newzealand part 1: J.Linn. Sci. Bot.vol. IV:333-357 .
- 33) **Chamberlaini, Y. M. 1965.** The marine algae of Gougn Island. Brit. MUS. (Naturlist.) Bot.vol.3(5):173-233.
- 34) **Coppeians, E. 1992.** Marine of Papua New Guinea (Madang Prov.) 2. A revised and completed list of Caulerpa (Caulerpa - Caulerpales). Blumea 36: 383 - 410 .
- 35) **Coppeians, E. & a. Marine of Papua New Guinea (Madang Prov.) Caulerpacoae (Caulerpa - Caulerpales) Blumea 33: 181 - 196 .**
- 36) **Coppeians, E. & W. F. Prud'homme Van Rein 1989.** Seaweeds of the snellius ll. Expedition Chlorophyta: Caulerpaes (except Caulerpa & Halimeda). Blumea Vol.34:119-124 .
- 37) **Dickinson 1956.** How to know seaweeds. U.S.A .
- 38) **Dickinson, C.I.&V.J. Foote 1951.** Marine algae from the Gold coast ll. kew, Bull.133-139 .
- 39) **El-Mohsen, A. E. F. A & Khoja 1973.** Studies on certain algenophyts from the Arabian Gulf of Saudi Arabia. Bull. Faculty. Science Riyadh Univ. 5:71-86 .
- 40) **Endicher, S. L. & Diesing C. M. 1845.** Enumeratio algarum quas an oram insulae karek sinus persici,legit Theodorus kotscl. Botanische zuitung. 3:268-69 .
- 41) **Gabary, D. J., G.I.Habseb & R. F.Scageh 1980.** The marine algae of British Colomdia and Northern Washington.: Divesion Rhodophyta (Red algae) Class Banriophceae Syesis, Vol. 13:137-195 .
- 42) **Gabrielson, P. W & G. T. Kraft 1984.** The Marine algae of Lord Howe Island (N.S.W) The famiy Solieriaceae (Gigartinales) Brunonia vol.7:217-251
- 43) **Gabrielson, P. W & G. T. Agardhiella** versus Neogardhiella Versus Neogardhiella (Solirridaa Rhodophyta) : An other look at the lectotina tenera. Taxon 34(2):275-300 .
- 44) **Gargiulo, G. M , F. D. E Masi & Tripodi 1992.** Morphology, reproduction and Taxonomy of Mediterranean species of Gracilarial . (Gracilariales, Rhodophyta). Phycologia 31:53-80 .
- 45) **Godeh, M. M. M. Nizamuddin & F. A. EL-Menifi, 1992.** Marine algae from easern coast of Libya (Cyrenai-ca). Pak.J.Bot.24(1):11-22
- 46) **Haas-Niekerk, T.D. 1965.** The genus Sphacelaria Lyngbye (Phaeopnyceae) in Netherlands. Blumea. Vol. 13 (1):145-161 .
- 47) **Hatta, A. M & W. F. Prud'nomme Van Rein 1991.** A Taxonomic revision of Indonesian Gelidiales (Rhodophyta). Blumea Vol.35:374-380 .
- 48) **Heerebout, G. R. 1968.** Studies on the Erythropelticlaceae (Rhodophceae Banriopnyceae). Blumea, vol.16 (1):138-157 .
- 49) **Heiba, H. I. M. M. Dugham, S. A. AL- Naydy & A. M. Rizk. 1990.** Phytochemical studies on the marine

- algae of Qatar Aradian Gulf Autar Univ. Sci. BULL.10:99-113 .
- 50) **Huisman, J. M. 1986.** The red algal Genus *Scinaia* (Galaxauraceae, Nemaliales) from Australia. *Phycologia*, vol.25(3):271-296 .
- 51) **Islam, N. A. K. M. 1976.** Contribution to study of marine algae of Banglaesg 253pp. Vaduz. Germany .
- 52) **Jonansen, H. W. & H. B. S. Womersley 1994.** *Jania* (Coralinales Rhodophyta) in southern Australia, Aust. syst. Bot. Vol. 7: 605-625 .
- 53) **Jagtap, T. G. 1992.** Marine flora of Nicobar Group of Island in andaman sea. Indian sia. Indian. J.Mar. Sci, Vol.21:56-58 .
- 54) **Jonen, O. M. P. J. A. Pugh & I. Titly 1994.** Observations on the benthic marine algal flora of south georgia: A floristic and ecological analysis. Bull. Nat. Hist. Mus. lond. (Bot) vol.24(2): 101-114 .
- 55) **Jones, D. A. 1986.** A field guide to the seashores of kuwait and the Arabian Gulf Univ.Gulf. Univ. Kuwat. 192pp.
- 56) **Jones,P. L. & W. M. J. Woelkerling 1984.** An analysis of trichocyste ad spore germination attributes at taxonomic characteric the phaeophylltc-fosilla complex (Coralinaceae-Rhodophyta). *Phycologia* Vol.23:183-194.
- 57) **Kamel, B. S. 1981.** Chemical composition of Arabian Gulf seaweeds. Lebensmittelwissen schaftlich uber technologie 14:160-62.
- 58) **Kraft,G. T. & J. Olsen-Stojkovich 1985.** *Avrainvillea calthina* (Udoteaceae Bryopsidales), a new green alga from Lord Howo Island, N.S.W. Australia. *Phycologia* Vol.24(3); 339-345.
- 59) **Krishnmurthy, V. & H. V. Joshi 1969.** The Species of *Ulva* L. from Indian waters. *Bot. J.Soc. Vol.62:123-130.*
- 60) **Koppen,J.D. 1975.** A Morphological and taconmc considerafion of tabellaria (Bacillariohtceae) frome the central Vnites states. *J.phycology*. VoL.11:236-244.
- 61) **Lawson, G. W. & John 1987.** The Marine algae of tropical environment of et Africa .
- 62) **Lawson, G. G, R. H. Simons & W. D. Isace 1990.** The Marine algal flora of Namibia. its distribution and affinities. *Bull. Brit. Mus. Nat. Hist(Bot)*. vol.20:153-168.
- 63) **Levring, T. 1951.** The marine algae of Australin I. Rhodophyta: Goniotrichales Bangiales and Nemalion ales. *Arkov. for Botanik Bond 2, No 6:457-529.*
- 64) **Littler, D. S. & M. M. Littler 1992.** Systematics of Avrainvillea (Bryopsidales Chlorophyta) in the tropical western Atlantic. *Phycology*, Vol.31 (5):375-418.
- 65) **McCain, J. C. 1984.** Marine ecology of saudi Arabia, The near-shores soft-bottom benthic comunities of the northern area Arabian.
- 66) **Millar, A. J. & G. T. Kraft 1994.** Catalogue of Marine Brown algae (Phaeophyta) of New South Wales, including Lord-Howe Island,south western pacific. *Aust.J.Bot.Vol. 7:1-64.*
- 67) **Millar, A. J. & G. T. Kraft 1994.** Catalogue of Marine Benthic Green algae (Chloropnyta) New South

- Wales,including Lord-Howe Island, south western pacific. Aust.J.Bot.Vol 7:419-453.
- 68) **Millar, A. J. K. 1990.** Marine red algae of the Goffs Harbour region, Northern New South Wales. Aust.Bot,Vol.3:293-593.
- 69) **Moghadam, F. 1974.** The Blue-green algae of gene Hot spring(IRAN). Bull.facul.sci. Tehran.Univ. Vol. 6 (2):78-152.
- 70) **Mshigeni, K. E. & M. M. Durgham 1987.** Benthic marine algae of Qatar, A preliminary survey. Unesco Regional Office, Doha.
- 71) **Mozzam, M. & M. Shameel 1985.** Studies on Bangiophceae(Rhodophyta) from the coast of Karachi. Pak.J.Bot.Vol 17(1):141-152.
- 72) **Nizamuddin, M. & F. Gessner 1970.** The Marine algae of northern part of Arabian sea and of the persian Gulf (Meteor) Forschung Ergebnies Reine D. 6:1-42.
- 73) **N'yeurt, A. D. 1996.** A preliminary floristic survey of the Benthic marine algae of Rotuman Island. Aust .Syst.Bot,Vol.9:361-490.
- 74) **Papenfuss, G. F. 1968.** Contribntion on the Knowledge of the Red sea.A history, coatalogue and biblioghrphy of Red sea benthic algae. Israel. J.Bot.vol.17:1-118.
- 75) **Phillips, J. A. , R.J. King. T. Tanaka & A. Mostaer 1993.** Steochospermum (Dictyotales, phaeophyceae),a poorly known algal genus,newly recorded in Australia Phycol.vol.32 (5) : 395-398.
- 76) **Penrose, D. & Y. M. Chamberlain 1993.** Hydrolitnon farinasum(Lamx) comb.nov., impelication for generic concepts in the castophoreideae (Cor allinaceae, Rhodophyta). Phycologia, Vol.32:295-303.
- 77) **Price, I. R. & F. J. Scott 1992.** The turf Algal flora of the great barier reef.266pp. Quensland, Australia.
- 78) **Richardson, W. D. 1975.** The Marine Algae of Trinidad,west Indies. Bull. Brit.Mus. (Nat.Hist.),vol.5(3); 47-145.
- 79) **Ricker, R. W. 1987.** Taxonomy and Biogeography of Macquaria Island seaweeds. Brit.Mus. (Nat.Hist.), 344pp.
- 80) **Rao, M., M. V. Murty & D. Satyanarayana 1994.** Nutrient realationships in the seaweeds of Visaknptnam coast of India,Ind.J.Mar.sci,Vol. 23: 43-46.
- 81) **Rao,U. M. & T. Sreeremulu 1970.** An annotated list of the marine algae of Visakhaotanam (India). Bot. J. Linn. soc,Vol.63:23-45.
- 82) **Solms-Laudach, H.G. 1894.** Mono graph of the Acetabularia. Trans. Linn. soci. secend seris, Vol.V:1-42.
- 83) **Srinivasan, K. S. 1969.** Pheologia Indica(lcones of lndkian marine algae). vol.1 Botanical survey of lhdian Calkutta. 52pp.
- 84) **Steentoft, M. 1967.** A revision of the marine algae of saotcome and principe (Gulf of Guinea). J. Linn. Soc. (Bot.). vol. 60 (382) ; 99-146.
- 85) **Steentoft, M. L. M. Irvine & W. F. Farnham 1995.** Two terete species of Gracilaria and Grcilariopsis (Gracilariales,Rhodophyta) in Britan. Phycologia, 34:113-127.
- 86) **Santelices, B. & S. Montalava. 1983.** Taxonomic studies on Gelidiaceae

- (Rhodophya) from central Chile. *Phycologia*. vol.22(2):185-196.
- 87) **Tanner, C. E. 1986.** Investigation of the Taxonomy and Morphologic variation *ulva* (Chlorophyta), *Ulva californica* Will. *Phycologia*. Vol. 25:510-520.
- 88) **Taylor, W. R. 1966.** West Indian marine algae. *Contrib. Univ. Mich. Herb.* 127-203.
- 89) **Taylor, W. R. 1967.** Species of *Caulerpa*(Chlorophyceae) collected on the international Indian Indian ocean expedition J.Ecology. Vol.16:45-53.
- 90) **Taylor, W. R. 1957.** Marine algae of the northeastern coast of north America (second revised edition). Univ. Mich. U.S.A.
- 91) **Taylor, W. M. 1964.** The genus *Turbinaria* in eastern seas. *J.Linn.soc.(Bot)* vol.58(375):475-490.
- 92) **Taylor, W. R. 1960.** Marine algae of the eastern tropical and subtropical coasts of the Americas Univ. mich. studies. sci.
- 93) **Verheij, E. & W.F. Prud'homme van Rein 1993.** Seaweeds of the Spermonde Archipelago SW.Sulawesi Indonesia. *Blumea*. Vol.37:385-510
- 94) **Vandermeulen, H., R. E. Dewreede & K.M. Cole 1984.** Nomenclatural recommendation tor three species of *Colpomenia* (Scytoniphonales, phoe phyta). *Taxon*. Vol.33:324-329.
- 95) **Van Din Hoek & Flinterman 1968.** The life history of *Sphacelaria furcigera* Kutzning (Phaeophyceae). *Blumea*. Vol.xV.l(1):196-242.
- 96) **Withell, W. F. A. J. K. Millar & G.T. kraft 1994.** Taxonomic studies of the genus *Gracilaria* (Gracilariales, Rhodophyta) from Austuralia. *Aust. Sysl. Bot.* Vol.27:281-352.
- 97) **Wiks, K. M. & W. J. Woelkerling 1991.** Southern Australiinn species of *Melodesia* (Corallinaceae, Rhodophyta). *Phycologia* Vol. 30(6):507-533.
- 98) **Womersley, H. B. S. 1979.** Southern Australian species of *Polysiphonia* Greville (Rhodophyta). *Aust.Syst.Bot.* Vol.7:459-528.
- 99) **Wynne, M. J. 1995.** Benthic Marine algae from the Seymhelles collected durig the R/V Te Vega Indian ocean expedition. *Conti. Univ. Mich. Her.* Vol.20:261-346.
- 100) **Zahid, P. B. M. Begum & M. Nizamuddin 1981.** Contribution to marine algae of Pakistan Rhodophyceae, *Polysiphonia* Greville 1823. *Pak. J. Bot.* 13(2):195-220.