

DR. IDOIA BIURRUN (Orcid ID : 0000-0002-1454-0433)

DR. REMIGIUSZ PIELECH (Orcid ID : 0000-0001-8879-3305)

DR. IWONA DEMBICZ (Orcid ID : 0000-0002-6162-1519)

PROF. FRANÇOIS GILLET (Orcid ID : 0000-0002-3334-1069)

MR. ŁUKASZ KOZUB (Orcid ID : 0000-0002-6591-8045)

CORRADO MARCENÒ (Orcid ID : 0000-0003-4361-5200)

PROF. ROBIN PAKEMAN (Orcid ID : 0000-0001-6248-4133)

MRS. IRENA AXMANOVÁ (Orcid ID : 0000-0001-9440-7976)

DR. SABINA BURRASCANO (Orcid ID : 0000-0002-6537-3313)

DR. STEFFEN BOCH (Orcid ID : 0000-0003-2814-5343)

PROF. HANS HENRIK BRUUN (Orcid ID : 0000-0003-0674-2577)

DR. TIMO CONRADI (Orcid ID : 0000-0003-2360-9284)

DR. BORJA JIMÉNEZ-ALFARO (Orcid ID : 0000-0001-6601-9597)

DR. RICARDA PÄTSCH (Orcid ID : 0000-0002-3349-0910)

DR. MASSIMO - TERZI (Orcid ID : 0000-0001-8801-6733)

DR. MANUELA WINKLER (Orcid ID : 0000-0002-8655-9555)

DR. JIANSHUANG WU (Orcid ID : 0000-0002-6768-8255)

DR. JUHA ALATALO (Orcid ID : 0000-0001-5084-850X)

MR. KUBER PRASAD PRASAD BHATTA (Orcid ID : 0000-0001-7837-1395)

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/JVS.13050](https://doi.org/10.1111/JVS.13050)

This article is protected by copyright. All rights reserved

PROF. ALESSANDRO CHIARUCCI (Orcid ID : 0000-0003-1160-235X)

DR. CRISTINA CHOCARRO (Orcid ID : 0000-0003-0405-0737)

DR. MARTA CZARNOCKA-CIECIURA (Orcid ID : 0000-0002-0168-0410)

DR. PATRYK MICHAŁ CZORTEK (Orcid ID : 0000-0002-4909-8032)

PROF. BALÁZS DEÁK (Orcid ID : 0000-0001-6938-1997)

DR. MARTIN DIEKMANN (Orcid ID : 0000-0001-8482-0679)

DR. JIŘÍ DOLEŽAL (Orcid ID : 0000-0002-5829-4051)

PROF. HAMID EJTEHADI (Orcid ID : 0000-0003-0681-5377)

DR. MANFRED FINCKH (Orcid ID : 0000-0003-2186-0854)

DR. ITZIAR GARCIA-MIJANGOS (Orcid ID : 0000-0002-6642-7782)

DR. ROSARIO GAVILÁN (Orcid ID : 0000-0002-1022-445X)

DR. MONIKA JANIŠOVÁ (Orcid ID : 0000-0002-6445-0823)

DR. ZYGMUNT KAŹKI (Orcid ID : 0000-0002-2241-1631)

DR. JUTTA KAPFER (Orcid ID : 0000-0002-8077-8917)

MR. ANDRÁS KELEMEN (Orcid ID : 0000-0002-2480-5669)

DR. LORENZO LAZZARO (Orcid ID : 0000-0003-0514-0793)

DR. JAN LEPS (Orcid ID : 0000-0002-4822-7429)

DR. DIEGO LIENDO (Orcid ID : 0000-0002-4816-4093)

DR. ZDEŇKA LOSOSOVÁ (Orcid ID : 0000-0001-9152-7462)

DR. ARANTZAZU L. LUZURIAGA (Orcid ID : 0000-0001-5023-7813)

DR. SIMONA MACCHERINI (Orcid ID : 0000-0002-2025-7546)

DR. ELIANE SERAINA MEIER (Orcid ID : 0000-0002-9545-1167)

DR. MICHELE MUGNAI (Orcid ID : 0000-0003-4315-2920)

DR. ALIREZA NAQINEZHAD (Orcid ID : 0000-0002-4602-6279)

DR. ARKADIUSZ NOWAK (Orcid ID : 0000-0001-8638-0208)

DR. SALZA PALPURINA (Orcid ID : 0000-0003-0416-5622)

PROF. ROBERT K. PEET (Orcid ID : 0000-0003-2823-6587)

DR. EULÀLIA PLADEVALL-IZARD (Orcid ID : 0000-0002-6693-5314)

DR. ZUZANA PLESKOVÁ (Orcid ID : 0000-0003-0153-1623)

DR. MARIA PILAR RODRÍGUEZ-ROJO (Orcid ID : 0000-0001-5449-9386)

DR. SOLVITA RUSINA (Orcid ID : 0000-0002-9580-4110)

MR. ANVAR SANAEI (Orcid ID : 0000-0001-8334-6944)

MR. VASCO SILVA (Orcid ID : 0000-0003-2729-1824)

DR. JUDIT SONKOLY (Orcid ID : 0000-0002-4301-5240)

MS. MARTA GAIA SPERANDII (Orcid ID : 0000-0002-2507-5928)

DR. SEBASTIAN ŚWIERSZCZ (Orcid ID : 0000-0003-2035-0035)

MR. AMIR TALEBI (Orcid ID : 0000-0002-5065-1368)

DR. LUBOMÍR TICHÝ (Orcid ID : 0000-0001-8400-7741)

DR. CSABA TÖLGYESI (Orcid ID : 0000-0002-0770-2107)

DR. PÉTER TÖRÖK (Orcid ID : 0000-0002-4428-3327)

PROF. ORSOLYA VALKÓ (Orcid ID : 0000-0001-7919-6293)

DR. THOMAS VANNESTE (Orcid ID : 0000-0001-5296-917X)

DR. RISTO VIRTANEN (Orcid ID : 0000-0002-8295-8217)

MISS YUN WANG (Orcid ID : 0000-0002-8238-5367)

DR. DAVID ZELENY (Orcid ID : 0000-0001-5157-044X)

DR. LI-QING ZHAO (Orcid ID : 0000-0002-8478-5786)

DR. JÜRGEN DENGLER (Orcid ID : 0000-0003-3221-660X)

Article type : Special Feature Article

Holger Kreft

Research Article for the Special Feature on Macroecology of Vegetation

Benchmarking plant diversity of Palaeartic grasslands and other open habitats

Running title: Plant diversity of Palaeartic grasslands

Idoia Biurrun¹, Remigiusz Pielech^{2,3}, Iwona Dembicz^{4,5}, François Gillet⁶, Łukasz Kozub⁴, Corrado Marcenò^{1,7}, Triin Reitalu⁸, Koenraad Van Meerbeek⁹, Riccardo Guarino¹⁰, Milan Chytrý⁷, Robin J. Pakeman¹¹, Zdenka Preislerová⁷, Irena Axmanová⁷, Sabina Burrascano¹², Sándor Bartha¹³, Steffen Boch¹⁴, Hans Henrik Bruun¹⁵, Timo Conradi¹⁶, Pieter De Frenne¹⁷, Franz Essl¹⁸, Goffredo Filibeck¹⁹, Michal Hájek⁷, Borja Jiménez-Alfaro²⁰, Anna Kuzemko²¹, Zsolt Molnár¹³, Meelis Pärtel²², Ricarda Pätzsch²³, Honor C. Prentice²⁴, Jan Roleček²⁵, Laura M.E. Sutcliffe²⁶, Massimo Terzi²⁷, Manuela Winkler^{28,29}, Jianshuang Wu³⁰, Svetlana Aćić³¹, Alicia T.R. Acosta³², Elias Afif³³, Munemitsu

Akasaka³⁴, Juha M. Alatalo³⁵, Michele Aleffi³⁶, Alla Aleksanyan³⁷, Arshad Ali³⁸, Iva Apostolova³⁹, Parvaneh Ashouri⁴⁰, Zoltán Bátor⁴¹, Esther Baumann⁴², Thomas Becker⁴³, Elena Belonovskaya⁴⁴, José Luis Benito Alonso⁴⁵, Asun Berastegi⁴⁶, Ariel Bergamini¹⁴, Kuber Prasad Bhatta⁴⁷, Ilaria Bonini⁴⁸, Marc-Olivier Büchler⁵, Vasyl Budzhak⁴⁹, Álvaro Bueno⁵⁰, Fabrizio Buldrini⁵¹, Juan Antonio Campos¹, Laura Cancellieri¹⁹, Marta Carboni³², Tobias Ceulemans⁵², Alessandro Chiarucci⁵¹, Cristina Chocarro⁵³, Luisa Conti^{32,157,158}, Anna Mária Csergő⁵⁴, Beata Cykowska-Marzencka^{55,5}, Marta Czarniecka-Wiera^{56,57}, Marta Czarnocka-Cieciura⁵⁸, Patryk Czortek⁵⁹, Jiří Danihelka^{7,60}, Francesco de Bello⁶¹, Balázs Deák⁶², László Demeter⁶³, Lei Deng⁶⁴, Martin Diekmann⁶⁵, Jiri Dolezal^{66,60}, Christian Dolnik⁶⁷, Pavel Dřevojan⁷, Cecilia Dupré⁶⁵, Klaus Ecker¹⁴, Hamid Ejtehadi⁶⁸, Brigitta Erschbamer⁶⁹, Javier Etayo⁷⁰, Jonathan Etzold⁷¹, Tünde Farkas⁷², Mohammad Farzam⁷³, George Fayvush³⁷, María Rosa Fernández Calzado⁷⁴, Manfred Finckh⁷⁵, Wendy Fjellstad⁷⁶, Georgios Fotiadis⁷⁷, Daniel García-Magro¹, Itziar García-Mijangos¹, Rosario G. Gavilán⁷⁸, Markus Germany⁷⁹, Sahar Ghafari⁸⁰, Gian Pietro Giusso del Galdo⁸¹, John-Arvid Grytnes⁴⁷, Behlül Güler⁸², Alba Gutiérrez-Girón⁸³, Aveliina Helm²², Mercedes Herrera¹, Elisabeth M. Hüllbusch⁸⁴, Nele Ingerpuu²², Annika K. Jägerbrand⁸⁵, Ute Jandt^{86,87}, Monika Janišová⁸⁸, Philippe Jeanneret⁸⁹, Florian Jeltsch⁹⁰, Kai Jensen⁹¹, Anke Jentsch⁸⁴, Zygmunt Kącki⁵⁷, Kaoru Kakinuma⁹², Jutta Kapfer⁷⁶, Mansoureh Kargar⁹³, András Kelemen⁶², Kathrin Kiehl⁹⁴, Philipp Kirschner⁶⁹, Asuka Koyama⁹⁵, Nancy Langer⁹⁶, Lorenzo Lazzaro⁹⁷, Jan Lepš⁶⁶, Ching-Feng Li^{98*}, Frank Yonghong Li⁹⁹, Diego Liendo¹, Regina Lindborg¹⁰⁰, Swantje Löbel¹⁰¹, Angela Lomba¹⁰², Zdeňka Lososová⁷, Pavel Lustyk⁷, Arantzazu L. Luzuriaga¹⁰³, Wenhong Ma⁹⁹, Simona Maccherini⁴⁸, Martin Magnes¹⁰⁴, Marek Malicki^{105,106}, Michael Manthey¹⁰⁷, Constantin Mardari¹⁰⁸, Felix May¹⁰⁹, Helmut Mayrhofer¹⁰⁴, Eliane Seraina Meier⁸⁹, Farshid Memariani¹¹⁰, Kristina Merunková⁷, Ottar Michelsen¹¹¹, Joaquín Molero Mesa⁷⁴, Halime Moradi¹¹², Ivan Moysiyanenko¹¹³, Michele Mugnai⁹⁷, Alireza Naqinezhad¹¹⁴, Rayna Natcheva³⁹, Josep M. Ninot¹¹⁵, Marcin Nobis¹¹⁶, Jalil Noroozi¹¹⁷, Arkadiusz Nowak^{118,119}, Vladimir Onipchenko¹²⁰, Salza Palpurina^{121,39}, Harald Pauli^{28,29}, Hristo Pedashenko³⁹, Christian Pedersen⁷⁶, Robert K. Peet¹²², Aaron Pérez-Haase^{115,123}, Jan Peters¹²⁴, Nataša Pipenbaher¹²⁵, Chrisoula Pirini¹²⁶, Eulàlia Pladevall-Izard¹¹⁵, Zuzana Plesková⁷, Giovanna Potenza¹²⁷, Soroor Rahmanian⁶⁸, Maria Pilar Rodríguez-Rojo¹²⁸, Vladimir Ronkin¹²⁹, Leonardo Rosati¹²⁷, Eszter Ruprecht¹³⁰, Solvita Rusina¹³¹, Marko Sabovljević¹³², Anvar Sanaei¹³³, Ana M. Sánchez¹⁰³, Francesco Santi⁵¹, Galina Savchenko¹²⁹, Maria Teresa Sebastià¹³⁴, Dariia Shyriaieva²¹, Vasco

Silva¹³⁵, Sonja Škornik¹²⁵, Eva Šmerdová⁷, Judit Sonkoly^{136,137}, Marta Gaia Sperandii^{138,32}, Monika Staniaszek-Kik¹³⁹, Carly Stevens¹⁴⁰, Simon Stifter¹⁴¹, Sigrid Suchrow⁹¹, Grzegorz Swacha⁵⁷, Sebastian Świerszcz^{118,142}, Amir Talebi¹¹², Balázs Teleki¹³⁶, Lubomír Tichý⁷, Csaba Tölgyesi⁴¹, Marta Torca¹, Péter Török^{136,137}, Nadezda Tsarevskaya⁴⁴, Ioannis Tsiripidis¹²⁶, Ingrid Turisova¹⁴³, Atushi Ushimaru¹⁴⁴, Orsolya Valkó⁶², Carmen Van Mechelen¹⁴⁵, Thomas Vanneste¹⁷, Iuliia Vasheniak¹⁴⁶, Kiril Vassilev³⁹, Daniele Viciani⁹⁷, Luis Villar¹⁴⁷, Risto Virtanen¹⁴⁸, Ivana Vitasović-Kosić¹⁴⁹, András Vojtkó¹⁵⁰, Denys Vynokurov²¹, Emelie Waldén¹⁰⁰, Yun Wang¹⁵¹, Frank Weiser⁴², Lu Wen⁹⁹, Karsten Wesche^{151,87,156}, Hannah White¹⁵², Stefan Widmer⁵, Sebastian Wolfrum^{153,154}, Anna Wróbel¹¹⁶, Zuoqiang Yuan¹³³, David Zelený¹⁵⁵, Liqing Zhao⁹⁹, Jürgen Dengler^{5,16,87}

*Deceased 29 November 2019

ORCIDs

Idoia Biurrún: <https://orcid.org/0000-0002-1454-0433>

Remigiusz Pielech: <https://orcid.org/0000-0001-8879-3305>

Iwona Dembicz: <https://orcid.org/0000-0002-6162-1519>

François Gillet: <https://orcid.org/0000-0002-3334-1069>

Łukasz Kozub: <https://orcid.org/0000-0002-6591-8045>

Corrado Marcenò: <https://orcid.org/0000-0003-4361-5200>

Triin Reitalu: <https://orcid.org/0000-0002-6555-3066>

Koenraad Van Meerbeek: <https://orcid.org/0000-0002-9260-3815>

Riccardo Guarino: <https://orcid.org/0000-0003-0106-9416>

Milan Chytrý: <https://orcid.org/0000-0002-8122-3075>

Robin J. Pakeman: <https://orcid.org/0000-0001-6248-4133>

Irena Axmanová: <https://orcid.org/0000-0001-9440-7976>

Sabina Burrascano: <https://orcid.org/0000-0002-6537-3313>

Sándor Bartha: <https://orcid.org/0000-0001-6331-7521>

Steffen Boch: <https://orcid.org/0000-0003-2814-5343>

Hans Henrik Bruun: <https://orcid.org/0000-0003-0674-2577>

Timo Conradi: <https://orcid.org/0000-0003-2360-9284>

Pieter De Frenne: <https://orcid.org/0000-0002-8613-0943>

Franz Essl: <https://orcid.org/0000-0001-8253-2112>

Goffredo Filibeck: <https://orcid.org/0000-0002-4187-9467>

Michal Hájek: <https://orcid.org/0000-0002-5201-2682>

Borja Jiménez-Alfaro: <https://orcid.org/0000-0001-6601-9597>

Anna Kuzemko: <https://orcid.org/0000-0002-9425-2756>

Zsolt Molnár: <https://orcid.org/0000-0001-5454-4714>

Meelis Pärtel: <https://orcid.org/0000-0002-5874-0138>

Ricarda Pätsch: <https://orcid.org/0000-0002-3349-0910>

Honor C. Prentice: <https://orcid.org/0000-0003-2548-0525>

Jan Roleček: <https://orcid.org/0000-0002-0023-1111>

Massimo Terzi: <https://orcid.org/0000-0001-8801-6733>

Manuela Winkler: <https://orcid.org/0000-0002-8655-9555>

Jianshuang Wu: <https://orcid.org/0000-0002-6768-8255>

Svetlana Ačić: <https://orcid.org/0000-0001-6553-3797>

Alicia T.R. Acosta: <https://orcid.org/0000-0001-6572-3187>

Munemitsu Akasaka: <https://orcid.org/0000-0002-0143-0841>

Juha M. Alatalo: <https://orcid.org/0000-0001-5084-850X>

Michele Aleffi : <https://orcid.org/0000-0001-8087-9550>

Alla Aleksanyan: <https://orcid.org/0000-0003-4073-1812>

Arshad Ali: <https://orcid.org/0000-0001-9966-2917>

Iva Apostolova: <https://orcid.org/0000-0002-2701-175X>

Parvaneh Ashouri: <https://orcid.org/0000-0003-3094-2376>

Zoltán Bátor: <https://orcid.org/0000-0001-9915-5309>

Thomas Becker: <https://orcid.org/0000-0002-7942-5575>

Elena Belonovskaya: <https://orcid.org/0000-0002-8354-4606>

José Luis Benito Alonso: <https://orcid.org/0000-0003-1086-8834>

Asun Berastegi: <https://orcid.org/0000-0003-0456-3305>

Ariel Bergamini: <https://orcid.org/0000-0001-8816-1420>

Kuber Prasad Bhatta: <https://orcid.org/0000-0001-7837-1395>

Ilaria Bonini: <https://orcid.org/0000-0002-9090-2926>

Marc-Olivier Büchler: <https://orcid.org/0000-0002-8967-8146>

Vasyl Budzhak: <https://orcid.org/0000-0002-7754-6437>

Alvaro Bueno: <https://orcid.org/0000-0002-1000-0886>

Fabrizio Buldrini: <https://orcid.org/0000-0002-0376-6760>

Juan Antonio Campos: <https://orcid.org/0000-0001-5992-2753>

Laura Cancellieri: <https://orcid.org/0000-0002-0102-259X>

Marta Carboni: <https://orcid.org/0000-0002-9348-4758>

Tobias Ceulemans: <https://orcid.org/0000-0001-6861-4613>

Alessandro Chiarucci: <https://orcid.org/0000-0003-1160-235X>

Cristina Chocarro: <https://orcid.org/0000-0003-0405-0737>

Luisa Conti: <https://orcid.org/0000-0001-8047-1467>

Anna Mária Csergő: <https://orcid.org/0000-0003-3325-2995>

Beata Cykowska-Marzencka: <https://orcid.org/0000-0002-5468-4909>

Marta Czarniecka-Wiera: <https://orcid.org/0000-0003-3294-5853>

Marta Czarnocka-Cieciura: <https://orcid.org/0000-0002-0168-0410>

Patryk Czortek: <https://orcid.org/0000-0002-4909-8032>

Jiří Danihelka: <https://orcid.org/0000-0002-2640-7867>

Francesco de Bello: <https://orcid.org/0000-0001-9202-8198>

Balázs Deák: <https://orcid.org/0000-0001-6938-1997>

Martin Diekmann: <https://orcid.org/0000-0001-8482-0679>

Jiri Dolezal: <https://orcid.org/0000-0002-5829-4051>

Pavel Dřevojan: <https://orcid.org/0000-0003-0802-3509>

Cecilia Dupré: <https://orcid.org/0000-0002-0867-4304>

Klaus Ecker: <https://orcid.org/0000-0003-3388-1353>

Hamid Ejtehadi: <https://orcid.org/0000-0003-0681-5377>

Brigitta Erschbamer : <https://orcid.org/0000-0002-6792-1395>

Javier Etayo: <https://orcid.org/0000-0003-0392-0710>

Jonathan Etzold: <https://orcid.org/0000-0002-2574-4415>

Mohammad Farzam: <https://orcid.org/0000-0003-1947-0187>

George Fayvush: <https://orcid.org/0000-0002-9710-2200>

Manfred Finckh: <https://orcid.org/0000-0003-2186-0854>

Wendy Fjellstad: <https://orcid.org/0000-0002-2849-9785>

Georgios Fotiadis: <https://orcid.org/0000-0003-0970-1607>

Itziar García-Mijangos: <https://orcid.org/0000-0002-6642-7782>

Rosario G. Gavilán: <https://orcid.org/0000-0002-1022-445X>

Sahar Ghafari: <https://orcid.org/0000-0002-9686-801X>

Gian Pietro Giusso del Galdo: <https://orcid.org/0000-0003-4719-3711>

John-Arvid Grytnes: <https://orcid.org/0000-0002-6365-9676>

Behlül Güler: <https://orcid.org/0000-0003-2638-4340>

Alba Gutiérrez-Girón: <https://orcid.org/0000-0002-0988-3343>

Aveliina Helm: <https://orcid.org/0000-0003-2338-4564>

Mercedes Herrera: <https://orcid.org/0000-0003-2812-866>

Annika K. Jägerbrand: <https://orcid.org/0000-0001-8016-0124>

Ute Jandt: <https://orcid.org/0000-0002-3177-3669>

Monika Janišová: <https://orcid.org/0000-0002-6445-0823>

Philippe Jeanneret: <https://orcid.org/0000-0002-6715-4632>

Florian Jeltsch: <https://orcid.org/0000-0002-4670-6469>

Kai Jensen: <https://orcid.org/0000-0002-0543-070X>

Anke Jentsch: <https://orcid.org/0000-0002-2345-8300>

Kaoru Kakinuma: <https://orcid.org/0000-0003-4647-9582>

Jutta Kapfer: <https://orcid.org/0000-0002-8077-8917>

András Kelemen: <https://orcid.org/0000-0002-2480-5669>

Kathrin Kiehl: <https://orcid.org/0000-0003-3931-187X>

Philipp Kirschner: <https://orcid.org/0000-0002-9263-5112>

Asuka Koyama: <https://orcid.org/0000-0002-7564-4332>

Lorenzo Lazzaro: <https://orcid.org/0000-0003-0514-0793>

Jan Lepš: <https://orcid.org/0000-0002-4822-7429>

Frank Yonghong Li: <https://orcid.org/0000-0002-5137-8017>

Diego Liendo: <https://orcid.org/0000-0002-4816-4093>

Regina Lindborg: <https://orcid.org/0000-0001-7134-7974>

Swantje Löbel: <https://orcid.org/0000-0001-9975-263X>

Angela Lomba: <https://orcid.org/0000-0003-3748-8751>

Zdeňka Lososová: <https://orcid.org/0000-0001-9152-7462>

Pavel Lustyk: <https://orcid.org/0000-0002-2514-9691>

Arantzazu L. Luzuriaga: <https://orcid.org/0000-0001-5023-7813>

Simona Maccherini: <https://orcid.org/0000-0002-2025-7546>

Martin Magnes: <https://orcid.org/0000-0002-5453-5735>

Marek Malicki: <https://orcid.org/0000-0003-0517-3560>

Michael Manthey: <https://orcid.org/0000-0002-1314-6290>

Felix May: <https://orcid.org/0000-0002-1106-8188>

Helmut Mayrhofer: <https://orcid.org/0000-0002-9433-3847>

Eliane Seraina Meier: <https://orcid.org/0000-0002-9545-1167>

Farshid Memariani: <https://orcid.org/0000-0001-5478-1859>

Ottar Michelsen: <https://orcid.org/0000-0001-7141-1948>

Joaquín Molero Mesa: <https://orcid.org/0000-0003-3106-8414>

Halime Moradi: <https://orcid.org/0000-0002-3738-9377>

Ivan Moysiyanenko: <https://orcid.org/0000-0002-0689-6392>

Michele Mugnai: <https://orcid.org/0000-0003-4315-2920>

Alireza Naqinezhad: <https://orcid.org/0000-0002-4602-6279>

Josep M. Ninot: <https://orcid.org/0000-0002-3712-0810>

Marcin Nobis: <https://orcid.org/0000-0002-1594-2418>

Jalil Noroozi: <https://orcid.org/0000-0003-4124-2359>

Arkadiusz Nowak: <https://orcid.org/0000-0001-8638-0208>

Vladimir Onipchenko: <https://orcid.org/0000-0002-1626-1171>

Salza Palpurina: <https://orcid.org/0000-0003-0416-5622>

Harald Pauli: <https://orcid.org/0000-0002-9842-9934>

Christian Pedersen: <https://orcid.org/0000-0002-9203-1108>

Robert K. Peet: <https://orcid.org/0000-0003-2823-6587>

Aaron Pérez-Haase: <https://orcid.org/0000-0002-5974-7374>

Nataša Pipenbahr: <https://orcid.org/0000-0003-1494-0194>

Chrisoula Pirini: <https://orcid.org/0000-0002-4511-1658>

Eulàlia Pladevall-Izard: <https://orcid.org/0000-0002-6693-5314>

Zuzana Plesková: <https://orcid.org/0000-0003-0153-1623>

Giovanna Potenza: <https://orcid.org/0000-0003-0957-1937>

Soroor Rahmanian: <https://orcid.org/0000-0001-6417-9781>

Maria Pilar Rodríguez-Rojo: <https://orcid.org/0000-0001-5449-9386>

Vladimir Ronkin: <https://orcid.org/0000-0003-3080-4117>

Leonardo Rosati: <https://orcid.org/0000-0003-4247-6157>

Eszter Ruprecht: <https://orcid.org/0000-0003-0122-6282>

Solvita Rusina: <https://orcid.org/0000-0002-9580-4110>

Marko Sabovljević: <https://orcid.org/0000-0001-5809-0406>

Anvar Sanaei: <https://orcid.org/0000-0001-8334-6944>

Ana María Sánchez: <https://orcid.org/0000-0002-6220-3001>

Francesco Santi: <https://orcid.org/0000-0002-9391-0059>

Galina Savchenko: <https://orcid.org/0000-0001-9436-7871>

Dariia Shyriaieva: <https://orcid.org/0000-0001-7140-7201>

Vasco Silva: <https://orcid.org/0000-0003-2729-1824>

Sonja Škornik: <https://orcid.org/0000-0002-7385-5507>

Eva Šmerdová: <https://orcid.org/0000-0003-4589-6317>

Judit Sonkoly: <https://orcid.org/0000-0002-4301-5240>

Marta Gaia Sperandii: <https://orcid.org/0000-0002-2507-5928>

Monika Staniaszek-Kik: <https://orcid.org/0000-0001-7969-0312>

Carly Stevens: <https://orcid.org/0000-0002-2390-1763>

Simon Stifter: <https://orcid.org/0000-0001-5957-9473>

Grzegorz Swacha: <https://orcid.org/0000-0002-6380-2954>

Sebastian Świerszcz: <https://orcid.org/0000-0003-2035-0035>

Amir Talebi: <https://orcid.org/0000-0002-5065-1368>

Balázs Teleki: <https://orcid.org/0000-0002-2417-0413>

Lubomír Tichý: <https://orcid.org/0000-0001-8400-7741>

Csaba Tölgyesi: <https://orcid.org/0000-0002-0770-2107>

Marta Torca: <https://orcid.org/0000-0001-7008-819X>

Péter Török: <https://orcid.org/0000-0002-4428-3327>

Nadezda Tsarevskaya: <https://orcid.org/0000-0002-0720-751X>

Ioannis Tsiripidis: <https://orcid.org/0000-0001-9373-676X>

Ingrid Turisova: <https://orcid.org/0000-0002-2753-5246>

Atushi Ushimaru: <https://orcid.org/0000-0002-2051-1233>

Orsolya Valkó: <https://orcid.org/0000-0001-7919-6293>

Carmen Van Mechelen: <https://orcid.org/0000-0002-9801-8975>

Thomas Vanneste: <https://orcid.org/0000-0001-5296-917X>

Luliia Vasheniak: <https://orcid.org/0000-0003-1020-3007>

Kiril Vassilev: <https://orcid.org/0000-0003-4376-5575>

Daniele Viciani: <https://orcid.org/0000-0003-3422-5999>

Risto Virtanen: <https://orcid.org/0000-0002-8295-8217>

Ivana Vitasović-Kosić: <https://orcid.org/0000-0001-9372-5892>

Denys Vynokurov: <https://orcid.org/0000-0001-7003-6680>

Emelie Waldén: <https://orcid.org/0000-0002-9180-2994>

Yun Wang: <https://orcid.org/0000-0002-8238-5367>

Karsten Wesche: <https://orcid.org/0000-0002-0088-6492>

Hannah White: <https://orcid.org/0000-0002-6793-8613>

Stefan Widmer: <https://orcid.org/0000-0002-4920-5205>

Sebastian Wolfrum : <https://orcid.org/0000-0003-0123-6720>

Anna Wróbel: <https://orcid.org/0000-0002-6713-7867>

Zuoqiang Yuan: <https://orcid.org/0000-0001-9197-7076>

David Zelený: <https://orcid.org/0000-0001-5157-044X>

Liqing Zhao: <https://orcid.org/0000-0002-5207-2490>

Jürgen Dengler: <https://orcid.org/0000-0003-3221-660X>

Author's institutional affiliations

¹Department of Plant Biology and Ecology, University of the Basque Country UPV/EHU, Bilbao, Spain

²Department of Forest Biodiversity, Faculty of Forestry, University of Agriculture in Kraków, Kraków, Poland

³Foundation for Biodiversity Research, ul. Terenowa 4c/6, 52-231 Wrocław, Poland

⁴Department of Ecology and Environmental Conservation, Institute of Environmental Biology,
Faculty of Biology, University of Warsaw, Warsaw, Poland

⁵Vegetation Ecology Group, Institute of Natural Resource Sciences (IUNR), Zurich University of
Applied Sciences (ZHAW), Wädenswil, Switzerland

⁶UMR Chrono-environnement, Université Bourgogne Franche-Comté, Besançon, France

⁷Department of Botany and Zoology, Faculty of Science, Masaryk University, Brno, Czech Republic

⁸Institute of Geology, Tallinn University of Technology, Tallinn, Estonia

⁹Division of Forest, Nature and Landscape, Department of Earth and Environmental Sciences, KU
Leuven, Leuven, Belgium

¹⁰Dept. STEBICEF - Botanical Unit, Università degli studi di Palermo, Palermo, Italy

¹¹Ecological Sciences, The James Hutton Institute, Aberdeen, United Kingdom

¹²Department of Environmental Biology, Sapienza University of Rome, Rome, Italy

¹³Institute of Ecology and Botany, Centre for Ecological Research, Vácrátót, Hungary

¹⁴Biodiversity & Conservation Biology, WSL Swiss Federal Research Institute, Birmensdorf,
Switzerland

¹⁵Section for Ecology and Evolution, Department of Biology, University of Copenhagen,
Copenhagen, Denmark

¹⁶Plant Ecology, Bayreuth Center of Ecology and Environmental Research (BayCEER), University of
Bayreuth, Bayreuth, Germany

¹⁷Forest & Nature Lab., Ghent University, Gontrode, Belgium

¹⁸Division of Conservation Biology, Vegetation and Landscape Ecology, Department of Botany and
Biodiversity Research, University of Vienna, Vienna, Austria

¹⁹Department of Agricultural and Forest Sciences (DAFNE), University of Tuscia, Viterbo, Italy

²⁰Research Unit of Biodiversity (CSIC, UO, PA), Oviedo University, Mieres, Spain

²¹Geobotany and Ecology Department, M.G. Kholodny Institute of Botany NAS of Ukraine, Kyiv, Ukraine

²²Institute of Ecology and Earth Sciences, University of Tartu, Tartu, Estonia

²³Wagner Ecology Lab, Department of Biological Sciences, University of Alberta, Edmonton, AB, Canada

²⁴Department of Biology, Lund University, Lund, Sweden

²⁵Institute of Botany, Department of Paleoecology, Czech Academy of Sciences, Brno, Czech Republic

²⁶Plant Ecology and Ecosystem Research, Georg-August University of Göttingen, Göttingen, Germany

²⁷Institute of Biosciences and Bioresources (IBBR), Italian National Council of Research (CNR), Bari, Italy

²⁸GLORIA co-ordination, Institute for Interdisciplinary Mountain Research, Austrian Academy of Sciences, Vienna, Austria

²⁹GLORIA co-ordination, Department of Integrative Biology and Biodiversity Research, University of Natural Resources and Life Sciences, Vienna (BOKU), Vienna, Austria

³⁰Institute of Environment and Sustainable Development in Agriculture, Chinese Academy of Agricultural Sciences, Beijing, China

³¹Department of Botany, Faculty of Agriculture, University of Belgrade, Belgrade-Zemun, Serbia

³²Department of Science, Roma Tre University, Rome, Italy

³³Department of Organisms and Systems Biology, Polytechnic School of Mieres, Oviedo University, Mieres, Spain

³⁴Institute of Agriculture, Tokyo University of Agriculture and Technology, Tokyo, Japan

³⁵Environmental Science Center, Qatar University, Doha, Qatar

³⁶School of Biosciences and Veterinary Medicine, University of Camerino, Camerino, Italy

³⁷Department of Geobotany and Plant Eco-Physiology, Institute of Botany aft. A.L. Takhtajyan NAS RA, Yerevan, Armenia

³⁸Forest Ecology Research Group, College of Life Sciences, Hebei University, Baoding, Hebei, China ³⁹Institute of Biodiversity and Ecosystem Research, Bulgarian Academy of Sciences, Sofia, Bulgaria

⁴⁰Agricultural Research, Education and Extension Organization (AREEO), Research Institute of Forests & Rangelands, Tehran, Iran

⁴¹Department of Ecology, University of Szeged, Szeged, Hungary

⁴²Biogeography, Bayreuth Center of Ecology and Environmental Research (BayCEER), University of Bayreuth, Bayreuth, Germany

⁴³Geobotany, Regional and Environmental Sciences, University of Trier, Trier, Germany

⁴⁴Institute of Geography, Russian Academy of Sciences, Moscow, Russia

⁴⁵JOLUBE Consultor Botánico, Jaca, Spain

⁴⁶Department of Biodiversity, Environmental Management of Navarre, Pamplona-Iruña, Spain

⁴⁷Department of Biological Sciences, University of Bergen, Bergen, Norway

⁴⁸Department of Life Sciences, University of Siena, Siena, Italy

⁴⁹Botany, Forestry and Horticulture, Yuriy Fedkovych Chernivtsi National University, Chernivtsy, Ukraine

⁵⁰Instituto de Recursos Naturales y Ordenación del Territorio (INDUROT), Universidad de Oviedo, Mieres, Spain

⁵¹Department of Biological, Geological and Environmental Sciences (BiGeA), University of Bologna, Bologna, Italy

⁵²Plant Conservation and Population Biology, Department of Biology, Leuven University, Leuven, Belgium

⁵³Department of Crop and Forest Science, Universitat de Lleida, Lleida, Spain

⁵⁴Department of Botany, Hungarian University of Agriculture and Life Sciences, Budapest, Hungary

⁵⁵Department of Mycology, W. Szafer Institute of Botany, Polish Academy of Sciences, Kraków, Poland

⁵⁶Department of Nature Protection and Rural Landscape, Institute of Technology and Life Sciences, Falenty, Poland

⁵⁷Botanical Garden, University of Wrocław, Wrocław, Poland

⁵⁸Laboratory of Statistical Analysis, National Information Processing Institute, Warsaw, Poland

⁵⁹Białowieża Geobotanical Station, Faculty of Biology, University of Warsaw, Białowieża, Poland

⁶⁰Institute of Botany, Czech Academy of Sciences, Průhonice, Czech Republic

⁶¹Desertification Research Centre (CIDE), Universitat de València & CSIC, Moncada, Spain

⁶²Lendület Seed Ecology Research Group, Centre for Ecological Research, Institute of Ecology and Botany, Vácrátót, Hungary

⁶³Department of Harghita County, National Agency for Protected Areas, Miercurea-Ciuc, Romania

⁶⁴Institute of Soil and Water Conservation, State Key Laboratory of Soil Erosion and Dryland Farming on the Loess Plateau, Northwest A&F University, Yangling, China

⁶⁵Vegetation Ecology and Conservation Biology, Institute of Ecology, FB 2, University of Bremen, Bremen, Germany

⁶⁶Department of Botany, Faculty of Science, University of South Bohemia, Ceske Budejovice, Czech Republic

⁶⁷Department of Landscape Ecology, Institute for Natural Resource Conservation, Kiel University, Kiel, Germany

⁶⁸Department of Biology, Ferdowsi University of Mashhad, Mashhad, Iran

⁶⁹Department of Botany, University of Innsbruck, Innsbruck, Austria

⁷⁰Pamplona, Spain

⁷¹ESTOK UG, Bernau (bei Berlin), Germany

⁷²Aggtelek National Park Directorate, Jósvalfó, Hungary

⁷³Department of Range and Watershed Management, Ferdowsi University of Mashhad, Mashhad, Iran

⁷⁴Department of Botany, Faculty of Pharmacy, University of Granada, Granada, Spain

⁷⁵Biodiversity, Evolution and Ecology of Plants (BEE), Institute of Plant Science and Microbiology (IPM), University of Hamburg, Hamburg, Germany

⁷⁶Department of Landscape Monitoring, Norwegian Institute of Bioeconomy Research, Ås, Norway

⁷⁷Department of Forestry and Natural Environment Management, Agricultural University of Athens, Karpenisi, Greece

⁷⁸Departament of Pharmacology, Pharmacognosy and Botany, Pharmacy, Complutense University, Madrid, Spain

⁷⁹Institute for Ecosystem Research, Kiel University, Kiel, Germany

⁸⁰Department of Range and Watershed Management, University of Mohaghegh Ardabili, Ardabil, Iran

⁸¹Department of Biological, Geological and Environmental Sciences, University of Catania, Catania, Italy

⁸²Biology Education, Dokuz Eylul University, Buca, İzmir, Turkey

⁸³Department of Biodiversity, Ecology and Evolution, Complutense University, Madrid, Spain

⁸⁴Disturbance Ecology, Bayreuth Center of Ecology and Environmental Research (BayCEER), University of Bayreuth, Bayreuth, Germany

⁸⁵Department of Environmental and Bioscience, the Rydberg Laboratory of Applied Science, Halmstad University, Halmstad, Sweden

⁸⁶Geobotany and Botanical Garden, Institute of Biology, Martin Luther University Halle-Wittenberg, Halle (Saale), Germany

⁸⁷German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, Leipzig, Germany

⁸⁸Institute of Botany, Plant Science and Biodiversity Center, Slovak Academy of Sciences, Banská Bystrica, Slovakia

⁸⁹Research Division Agroecology and Environment, Agroscope, Zürich, Switzerland

⁹⁰Plant Ecology and Nature Conservation, University of Potsdam, Potsdam, Germany

⁹¹Applied Plant Ecology, Biocentre Klein Flottbek, University of Hamburg, Hamburg, Germany

⁹²Asian Demographic Research Institute, Shanghai University, Shanghai, China

⁹³Natural Resources and Watershed Management Administration of Alborz Province, Karaj, Iran

⁹⁴Vegetation Ecology and Botany, Faculty of Agricultural Sciences and Landscape Architecture,
Osnabrück University of Applied Sciences, Osnabrück, Germany

⁹⁵Center for Biodiversity, Forestry and Forest Products Research Institute, Tsukuba, Japan

⁹⁶Stiftung Naturschutzfonds Brandenburg, Potsdam, Germany

⁹⁷Laboratory of Plant Systematics and Phytogeography, Department of Biology, University of
Florence, Florence, Italy

⁹⁸School of Forestry and Resource Conservation, National Taiwan University, Taipei, Taiwan
(deceased)

⁹⁹Ministry of Education Key Laboratory of Ecology and Resource Use of the Monoglian Plateau &
School of Ecology and Environment, Inner Mongolia University, Hohhot, China

¹⁰⁰Department of Physical Geography, Stockholm University, Stockholm, Sweden

¹⁰¹Landscape Ecology and Environmental Systems Analysis, Institute of Geoecology, TU
Braunschweig, Braunschweig, Germany

¹⁰²Research Centre in Biodiversity and Genetic Resources (CIBIO) - Research Network in
Biodiversity and Evolutionary Biology (InBIO), University of Porto, University of Porto, Vairão,
Vila do Conde, Portugal

¹⁰³Área de Biodiversidad y Conservación, Universidad Rey Juan Carlos, Móstoles, Spain

¹⁰⁴Institute of Biology, Department of Plant Sciences, University of Graz, Graz, Austria

¹⁰⁵Department of Botany, University of Wrocław, Wrocław, Poland

¹⁰⁶Botanical Garden of Medicinal Plants, Wrocław Medical University, Wrocław, Poland

¹⁰⁷Institute of Botany and Landscape Ecology, Greifswald University, Greifswald, Germany

¹⁰⁸Anastasie Fatu Botanical Garden, Alexandru Ioan Cuza University from Iași, Iași, Romania

¹⁰⁹Theoretical Ecology, Institute of Biology, Freie Universität Berlin, Berlin, Germany

¹¹⁰Department of Botany, Research Center for Plant Sciences, Ferdowsi University of Mashhad, Mashhad, Iran

¹¹¹Department of Industrial Economics and Technology Management, Norwegian University of Science and Technology, Trondheim, Norway

¹¹²Department of Plant Science, School of Biology, College of Science, University of Tehran, Tehran, Iran

¹¹³Department of Botany, Kherson State University, Kherson, Ukraine

¹¹⁴Department of Plant Biology, Faculty of Basic Sciences, University of Mazandaran, Babolsar, Iran

¹¹⁵Department of Evolutionary Biology, Ecology and Environmental Sciences, Faculty of Biology, University of Barcelona, Barcelona, Spain

¹¹⁶Institute of Botany, Jagiellonian University, Kraków, Poland

¹¹⁷Department of Botany and Biodiversity Research, University of Vienna, Vienna, Austria

¹¹⁸Botanical Garden Center for Biological Diversity Conservation in Powsin, Polish Academy of Sciences, Warsaw, Poland

¹¹⁹Institute of Biology, University of Opole, Opole, Poland

¹²⁰Ecology and Plant Geography, Lomonosov Moscow State University, Moscow, Russia

¹²¹National Museum of Natural History, Bulgarian Academy of Sciences, Sofia, Bulgaria

¹²²Department of Biology, University of North Carolina, Chapel Hill, NC, United States

¹²³Department of Biosciences, Faculty of Sciences and Technology, University of Vic - Central University of Catalonia, Vic, Spain

¹²⁴Michael-Succow-Foundation, Greifswald, Germany

¹²⁵Biology Department, Faculty of Natural Sciences and Mathematics, University of Maribor, Maribor, Slovenia

¹²⁶Department of Botany, School of Biology, Aristotle University of Thessaloniki, Thessaloniki, Greece

¹²⁷School of Agricultural, Forest, Food and Environmental Sciences, University of Basilicata, Potenza, Italy

¹²⁸Institute of Environmental Sciences, University of Castilla-La Mancha, Toledo, Spain

¹²⁹Department of Zoology and Animal Ecology, V.N. Karazin Kharkiv National University, Kharkiv, Ukraine

¹³⁰Hungarian Department of Biology and Ecology, Babeş-Bolyai University, Cluj-Napoca, Romania

¹³¹Faculty of Geography and Earth Sciences, University of Latvia, Riga, Latvia

¹³²Institute for Botany and Botanical Garden "Jevremovac", Faculty of Biology, University of Belgrade, Belgrade, Serbia

¹³³CAS Key Laboratory of Forest Ecology and Management, Institute of Applied Ecology, Chinese Academy of Sciences, Shenyang, China

¹³⁴Group GAMES and Department of Horticulture, Botany and Landscaping, University of Lleida, Lleida, Spain

¹³⁵Centre for Applied Ecology "Professor Baeta Neves", CEABN/InBIO, School of Agriculture, University of Lisbon, Lisbon, Portugal

¹³⁶Department of Ecology, University of Debrecen, Debrecen, Hungary

¹³⁷MTA-DE Lendület Functional and Restoration Ecology Research Group, Debrecen, Hungary

¹³⁸Department of Ecology, Centro de Investigaciones Sobre Desertificación (CSIC-UV-GV),
Moncada (Valencia), Spain

¹³⁹Department of Geobotany and Plant Ecology, University of Łódź, Łódź, Poland

¹⁴⁰Lancaster Environment Centre, Lancaster University, Lancaster, United Kingdom

¹⁴¹Institute for Alpine Environment, Eurac Research, Bolzano, Italy

¹⁴²The Franciszek Górski Institute of Plant Physiology, Polish Academy of Sciences, Kraków,
Poland

¹⁴³Faculty of Natural Sciences, Matej Bel University in Banská Bystrica, Banská Bystrica, Slovakia

¹⁴⁴Graduate School of Human Development and Environment, Kobe University, Kobe, Japan

¹⁴⁵PXL Bio-Research, PXL University of Applied Sciences and Arts, Diepenbeek, Belgium

¹⁴⁶Department of Chemistry, Biology and Biotechnologies, Vasyl' Stus Donetsk National
University, Vinnytsia, Ukraine

¹⁴⁷Instituto Pirenaico de Ecología IPE-CSIC, Consejo Superior de Investigaciones Científicas, Jaca
(Huesca), Spain

¹⁴⁸Ecology & Genetics, University of Oulu, Oulu, Finland

¹⁴⁹Faculty of Agriculture, Department of Agricultural Botany, University of Zagreb, Zagreb, Croatia

¹⁵⁰Department of Botany, Eszterházy Károly University of Applied Sciences, Eger, Hungary

¹⁵¹Department of Botany, Senckenberg Museum of Natural History Görlitz, Görlitz, Germany

¹⁵²School of Natural Sciences, Trinity College Dublin, The University of Dublin, Dublin, Ireland

¹⁵³Life Science Center Weihenstephan, Technical University of Munich, Freising, Germany

¹⁵⁴Institute for Organic Farming, Soil and Resource Management, Bavarian State Research Center for Agriculture, Freising, Germany

¹⁵⁵Institute of Ecology and Evolutionary Biology, National Taiwan University, Taipei, Taiwan

¹⁵⁶Technische Universität Dresden, International Institute Zittau, Zittau, Germany

¹⁵⁷Faculty of Environmental Sciences, University of Life Sciences Prague, Praha-Suchdol, Czech Republic

¹⁵⁸Institute of Botany, Czech Academy of Sciences, Třeboň, Czech Republic

Correspondence

Idoia Biurrun, Department of Plant Biology and Ecology, University of the Basque Country UPV/EHU, Bilbao, Spain

Email: idoia.biurrun@ehu.es

Funding Information

Funding information is provided in Appendix S7.

Abstract

Aims: Understanding fine-grain diversity patterns across large spatial extents is fundamental for macroecological research and biodiversity conservation. Using the GrassPlot database, we provide benchmarks of fine-grain richness values of Palaeartic open habitats for vascular plants, bryophytes, lichens and complete vegetation (i.e., the sum of the former three groups).

Location: Palaeartic biogeographic realm.

Methods: We used 126,524 plots of eight standard grain sizes from the GrassPlot database: 0.0001 m², 0.001 m², 0.01 m², 0.1 m², 1 m², 10 m², 100 m² and 1000 m² and calculated the mean richness and standard deviations, as well as maximum, minimum, median, and first and third quartiles for each combination of grain size, taxonomic group, biome, region, vegetation type and phytosociological class.

Results: Patterns of plant diversity in vegetation types and biomes differ across grain sizes and taxonomic groups. Overall, secondary (mostly semi-natural) grasslands and natural grasslands are the richest vegetation type. The open-access file "GrassPlot Diversity Benchmarks" and the web tool "GrassPlot Diversity Explorer" are now available online (<https://edgg.org/databases/GrasslandDiversityExplorer>) and provide more insights into species richness patterns in the Palaearctic open habitats.

Conclusions: The GrassPlot Diversity Benchmarks provide high-quality data on species richness in open habitat types across the Palaearctic. These benchmark data can be used in vegetation ecology, macroecology, , biodiversity conservation and data quality checking. While the amount of data in the underlying GrassPlot database and their spatial coverage are smaller than in other extensive vegetation-plot databases, species recordings in GrassPlot are on average more complete, making it a valuable complementary data source in macroecology.

Keywords: benchmark, bryophyte, fine-grain biodiversity, grassland, GrassPlot Diversity Explorer, lichen, open habitat, Palaearctic, scale dependence, species-area relationship, vascular plant, vegetation plot.

Main Text

Introduction

Documenting and understanding patterns of biodiversity is a central issue in biogeography and macroecology (Gaston, 2000; Barthlott et al., 2007; Pärtel et al., 2016) and is also fundamental for sustainable land use and biodiversity conservation (Whittaker et al., 2005), as ecosystem function and stability are dependent on biodiversity (Tilman & Downing, 1994; Hooper et al., 2005). The increasing awareness of the current environmental crisis makes biodiversity studies even more valuable and necessary, especially for ecosystems such as grasslands, which are massively threatened by land-use change (Fischer et al., 2018). Plant species richness has been mapped globally using coarse-grain data (Barthlott et al., 2005; Kier et al., 2005; Brummit et al., 2020). However, fine-grain data on the local co-occurrence of species in plant communities across continental or global spatial extents are required for macroecological studies that link diversity patterns and assembly processes (Bruehlheide et al., 2019). Nevertheless, information on broad-scale, fine-grain plant distribution is still scattered, inconsistent, and often of uncertain data quality, especially for bryophytes and lichens (Beck et al., 2012).

Vegetation plots stored in large databases (e.g., European Vegetation Archive, EVA, Chytrý et al., 2016; sPlot, Bruehlheide et al., 2019) are increasingly used to explore fine-grain plant diversity patterns and underlying assembly processes at continental to global extents (Bruehlheide et al., 2018; Večeřa et al., 2019; Axmanová et al., 2021). The use of vegetation plots also allows the identification of regions with high fine-grain α -diversity for particular habitats, called richness hotspots (Divíšek & Chytrý, 2018; Večeřa et al., 2019). However, it should be considered that vegetation plots derived from phytosociological sampling may vary in plot size by several orders of magnitude, even within the same vegetation type (Chytrý, 2001). Sometimes information on plot size may be lacking or only approximate. Therefore, diversity inference from phytosociological data has to consider plot sizes and should be interpreted with caution (Chytrý, 2001; Chytrý & Otýpková, 2003).

Ecologists and conservationists need reliable species richness benchmarks (i.e., maximum, minimum, mean and other basic statistics) to assess plant communities as being above or below

average in richness for a specific region or vegetation type (Yen et al., 2019). To produce reliable benchmarks, plot size should be integrated into any analysis, and large amounts of high-quality vegetation-plot data are needed. Previous studies providing global richness data at several plot sizes focused on maximum values and left out information on the distribution of richness values (Wilson et al., 2012; Chytrý et al., 2015). This information is needed for both fundamental research and biodiversity conservation (Dengler et al., 2016a; Yen et al., 2019), e.g., when establishing thresholds between average and species-rich grasslands or identifying species-poor degraded grasslands for restoration.

Palaeartic grasslands host a considerable part of the realm's diversity (Dengler et al., 2020a). At fine spatial grains (< 100 m²), they can even hold higher plant diversity than tropical forests (Wilson et al., 2012). After an early and rudimentary attempt of benchmarking grassland diversity globally (Faber-Langendoen & Josse, 2010), Dengler et al. (2016a) provided a first overview of benchmarking plant diversity of Palaeartic grasslands based on a relatively small dataset.

Here, we use GrassPlot, the database of multi-scale plant diversity in Palaeartic grasslands (<https://edgg.org/databases/GrassPlot>; Dengler et al., 2018), to provide comprehensive benchmarks of fine-grain plant richness values of Palaeartic open habitats across biomes at eight plot sizes, each separated by order of magnitude: 0.0001 m², 0.001 m², 0.01 m², 0.1 m², 1 m², 10 m², 100 m² and 1000 m². We aim to display hot- and cold-spots of fine-grain α -diversity (species richness) across biomes and vegetation types. Besides total plant richness (complete vegetation), we separately assess vascular plant, bryophyte and lichen richness, as it has been already demonstrated that the richness of these taxonomic groups should be assessed separately (Dengler *et al.*, 2016a). In summary, we (i) present major diversity patterns in Palaeartic open habitats that can be derived from GrassPlot, (ii) introduce the GrassPlot Diversity Benchmarks (a dataset made public together with this article) and the GrassPlot Diversity Explorer (an online tool released together with this article), and (iii) outline some potential applications and impacts of both.

Methods

Data compilation

We used plot-based data from the collaborative vegetation database GrassPlot (Dengler et al., 2018; <https://edgg.org/databases/GrassPlot>), affiliated to the Eurasian Dry Grassland Group (EDGG), and registered in the Global Index of Vegetation-Plot Databases (EU-00-003; GIVD; Dengler et al., 2011). GrassPlot incorporates standardized vegetation-plot data sampled in precisely delimited plots together with methodological, environmental and structural information from open habitats, e.g. grasslands and other plant communities dominated by herbs, dwarf-shrubs or cryptogams from the Palaeartic biogeographic realm (i.e., Europe, North Africa, and West, Central, North and East Asia). Besides individual plots, GrassPlot specifically contains multi-scale datasets from nested-plot sampling schemes (e.g., Dengler et al., 2016b) with plot sizes ranging from 0.0001 to 1,024 m². The last published version of the database (GrassPlot v.2.00; Biurrun et al., 2019) contained more than 190,000 plots of different grain sizes across 22,422 individual plots and 5,749 nested-plot series with at least two grain sizes.

For this benchmarking study, we retrieved all plots with grain sizes 0.0001 m², 0.001 m², 0.01 m², 0.1 m², 1 m², 10 m², 100 m² and 1000 m² contained in GrassPlot v.2.10 (version of 1 Oct 2020), belonging to 225 datasets (Appendix S1). According to the typical species-area relationships in Palaeartic grasslands (Dengler et al., 2020b), 10% difference in the area means only about 2% difference in richness or less, which is negligible compared to any other source of richness variation. Thus, 2,372 plots deviating less than 10% from standard grain sizes (0.0009 m², 0.09 m², 9 m², 10.89 m², 900 m² and 1024 m²) were also selected and used for the benchmarks of the respective grain size. The final dataset contained 126,524 plots (Table 1) distributed across 49 countries (Appendix S2, Table S2.1), eight biomes throughout the Palaeartic realm and a wide range of open habitats, such as grasslands, dwarf shrublands, wetlands, and deserts (Figure 1, Table S2.2).

Biomes were assigned using the biome classification provided in Bruehlheide et al. (2019), which is based on the nine ecozones of Schultz (2005) plus an additional alpine biome based on Körner et al. (2017). Plots were also assigned to ten geographic regions following Dengler et al. (2020a). We created a two-level vegetation typology with 22 vegetation types grouped into six coarse

categories: natural grasslands, secondary grasslands, azonal communities, dwarf shrublands, tall-forb and ruderal communities and deserts and semi-deserts (more details in Appendix S2, Table S2.2 and Figure S2.1). Plots were assigned to vegetation types based on expert knowledge either individually by data owners or using general assignment rules of phytosociological syntaxa to vegetation types (see Table S2.3). 75% of plots in the dataset have a phytosociological assignment at least at the class level.

GrassPlot includes plot data sampled following two alternative methods for recording the presence of vascular plant species: “rooted presence”, which only records individuals as present in the plot if they root inside, and “shoot presence”, which records individuals as present if any part of stems or leaves are inside the plot (Dengler, 2008). The majority of plots in the dataset were recorded using the “shoot presence” method, and 13.4% of plots used “rooted presence”, while only a small fraction (0.1%) used a combined method, where shrubs were recorded using “rooted presence” and grasses and forbs using the “shoot presence”, or the recording method was not known (0.2%).

For linguistic convenience, we include lichens under the generic term “plants”. Thus, we considered four taxonomic groups: vascular plants, bryophytes, lichens and complete vegetation (i.e., the sum of the former three groups).

Establishing and providing benchmark values

We calculated mean species richness values and standard deviations, as well as maximum, minimum, median, and first and third quartiles for each combination of grain size, taxonomic group, biome, region, country, vegetation type (at coarse and fine classification level), phytosociological class and method (shoot vs. rooted, nested series with seven standard grain sizes vs. any plots). The data are organized as a spreadsheet, in which each of the 728,396 lines represents one combination of these factors, and the columns provide the statistics, i.e., number of plots, number of independent observations, minimum, maximum, mean, standard deviation, median, and first and third quartiles. We call these data the GrassPlot Diversity Benchmarks and provide them in Appendix S3 in Supporting Information as a spreadsheet file (70 MB). This file is open access and is also provided on the website of the GrassPlot Diversity Explorer

(<https://edgg.org/databases/GrasslandDiversityExplorer>) for free download. We intend to update it at regular intervals while keeping former versions available to make any studies based on these data reproducible.

Many nested series contain several subplots of the same size. Sometimes these are multiple contiguous subplots covering the entire surface of the largest plot. Because of a high degree of spatial pseudoreplication, using these richness values separately for calculating mean richness might bias the results. Thus, for all benchmarks, except for the maximum and minimum richness, we used the averaged values of each grain size in each nested series, i.e., only the independent observations. The number of independent observations decreased from 126,524 to 48,449 plots (Table 1), 6,509 of them belonging to nested series with at least seven of our standard grain sizes, 16,499 belonging to nested series with less than seven standard sizes, and 25,441 individual plots. In the dataset containing only independent observations, the percentage of plots using “rooted presence” rose from 13.4 to 23.4%.

We also added two filtering options as they can have significant effects on resulting richness patterns. (1) We allow filtering for data that were sampled with “rooted presence” or “shoot presence”. As has been shown theoretically (Williamson, 2003) and empirically (Güler et al., 2016; Cancellieri et al., 2017; Zhang et al., 2021), species richness recorded with the rooted method deviates increasingly negatively from values recorded with the shoot-presence method as grain size decreases. (2) Subsetting to only those plots belonging to nested series with at least the seven “EDGG standard grain sizes” (0.0001 to 100 m²; see Dengler et al., 2016b) is also possible. This function can be important when analyzing species-area relationships, which otherwise might be distorted by uneven representation of different grain sizes in specific regions..

Richness hotspots

In this study we aim at identifying fine-grain α -diversity hotspots (hereafter, richness hotspots). These richness hotspots are different from biodiversity hotspot of Myers et al. (2000), who emphasized a concentration of endemic species in larger regions combined with severe habitat loss. Other criteria such as the number of rare or threatened species and total species richness

are also currently used to identify these hotspots; moreover, this term is now most commonly used with reference to regions of high species richness (Reid, 1998). Another difference with the most widely used concept of the biodiversity hotspot is that we are using fine-grain resolution (plot level, e.g., lower than 1 km²), while most studies identify hotspots using coarse-grain resolution maps, generally at 10,000 km² (Küper et al., 2004) or even coarser (Myers et al., 2000).

Development of the GrassPlot Diversity Explorer

The GrassPlot Diversity Explorer (<https://edgg.org/databases/GrasslandDiversityExplorer>) was developed to provide a dynamic version of the GrassPlot Diversity Benchmarks. We did this in R version 4.0.2 (R Core Team, 2020), using the ‘shiny’ package (Chang et al., 2020). We also used other R packages, including ‘tidyr’ and ‘dplyr’ for data preparation (Wickham & Henry, 2020; Wickham et al., 2020), ‘ggplot2’, ‘ggpubr’ and ‘sunburstsR’ for visualisation of the outcomes (Wickham, 2016; Bostock et al., 2020; Kassambara, 2020), ‘summarytools’ for generating summary statistics (Comtois, 2020), ‘leaflet’ for producing an interactive map (Cheng et al., 2019), and ‘shinyWidgets’ and ‘shinycssloaders’ to increase the functionality of the shiny package (Perrier et al., 2020; Sali & Attali, 2020). The GrassPlot Diversity Explorer was then deployed on a dedicated server using the ‘rsconnect’ package (Allaire, 2019).

Results

GrassPlot Diversity Benchmarks

Richness hotspots of vascular plants in grasslands and other open habitats are scattered across the Palaearctic. However, they may vary across grain sizes, both regarding mean richness (Figure 2) and maximum richness (Appendix S4, Figure S4.3). Richness hotspots also change according to vegetation type (Figure S4.1) and taxonomic group (Figure S4.2). Maximum richness hotspots of bryophytes, lichens and complete vegetation also vary with grain size (Figures S4.4 to S4.6).

Patterns of plant diversity in vegetation types differ across grain sizes (Figure 3) and taxonomic groups (Figure 4). Secondary grasslands show the highest mean richness of vascular plants across the smallest sizes, but natural grasslands are equally diverse in the largest ones (Figure 3); this

pattern is found in both the plots recorded using “rooted presence” and “shoot presence” if they are analyzed separately (Table S5.2). Plant diversity patterns in biomes are even more dependent on grain size, with the highest means in the Boreal biome at 0.01 and 10 m² (Figure 3). More detailed results on richness patterns across grain sizes for the combination of biomes and coarse-level vegetation types are shown in Appendix S5 (Figures S5.1 to S5.4 for complete vegetation, vascular plants, bryophytes, and lichens, respectively). This inconsistency of diversity patterns across grain sizes and taxonomic groups is even more evident if we use a fine-level vegetation typology (Figures S5.5 to S5.12), although some patterns emerge, such as meso-xeric grasslands showing the highest mean richness for vascular plants across most grain sizes. The mean richness of bryophytes, lichens and complete vegetation strongly vary with grain size. In addition to arctic-alpine heathlands, sandy dry grasslands, rocky grasslands and mesic grasslands show the highest values, as well as several azonal communities such as saline, rocks and screes, and wetlands (Figures S5.5 to S5.12). Maximum richness corresponds to secondary grasslands across most grain sizes, but once again, the pattern changes for bryophytes and lichens, with maxima often in natural grasslands (Table 2). As regards biomes, the maximum richness slightly changes across grain sizes and taxonomic groups, although the Temperate mid-latitudes hold most of the maxima for all taxonomic groups (Appendix S5, Table S5.1).

Species-area relationships (SARs) of the six best-represented grassland types show similar patterns for vascular plants and complete vegetation, both with a continuous upward curvature in the semi-log space (Figure 5). Meso-xeric grasslands show the highest means across grain sizes, both for vascular plants and for complete vegetation, while sandy dry grasslands are the poorest type. Alpine grasslands outperform meso-xeric grasslands for complete vegetation at smallest sizes due to the strong richness increase related to the inclusion of non-vascular plants. The relative importance of this fraction in the total richness is even stronger for sandy dry grasslands and Mediterranean grasslands, the latter reaching the second position in the richness ranking for the largest sizes. Figures S5.13 to S5.16 combine SARs generated with all GrassPlot data with SARs corresponding to the subset of plots belonging to nested series with seven standard grain sizes for all fine-level vegetation types and the four taxonomic groups. In most cases lines of SARs using all plots are below lines of SARs of the subset of nested series.

GrassPlot Diversity Explorer

The GrassPlot Diversity Explorer is an easy-to-use online interactive tool that provides users flexibility in exploring and visualizing richness data collected in the GrassPlot database. The GrassPlot Diversity Explorer can be accessed via the EDGG website (<https://edgg.org/databases/GrasslandDiversityExplorer>). The tool is organized into eight panels (Figure 6). The first panel shows species richness in different vegetation types, and two vegetation typologies are used, including the two-level vegetation typology used in the GrassPlot database (see Appendix S2, Table S2.2) and phytosociological classes (for details, see Appendix S2, Table S2.3). Users can explore species richness by generating boxplots for eight standard size grains and selected taxonomic groups. The second panel presents species richness in geographical regions and biomes and the third one species-area relationships (SARs) for selected vegetation units. Details on the two-level vegetation typology can be found in the fourth panel, and the fifth panel presents descriptive statistics for the selected dataset. Users can generate these statistics for filtered datasets based on taxonomic group, vegetation types and region of interest. Besides graphs and descriptive statistics, users can also explore the distribution of plots based on selected criteria. Filtered plots are displayed on background maps in the “Map” panel. The seventh panel contains links to the main file of the GrassPlot Diversity Benchmarks. Finally, the last panel, “Information”, explains the concepts of biomes, geographical regions and vegetation typologies used in the GrassPlot Diversity Explorer.

Discussion

General diversity patterns

Fine-grain α -diversity patterns of Palaearctic open habitats are not consistent across grain sizes, as shown in Figure 2. While grain size as a determinant of species richness patterns has previously been reported in many coarse-grain studies (see Rahbek, 2005, for a review), here we demonstrate that it is also influential at the scale of ecological communities. Consequently, plant richness hotspots in Palaearctic open habitats are not necessarily consistent across scales, as already demonstrated for vertebrates (Shriner et al., 2006; McKerrow et al., 2018). Given that

GrassPlot data have still large geographic gaps, we refrain from comparing our emerging richness hotspots with those obtained from coarse-grain data (e.g., Barthlott et al., 2005, for vascular plants; Geffert et al., 2013, for bryophytes) and with other community-level patterns in grasslands emerging from fine-grain data (e.g., Wilson et al., 2012; Chytrý et al., 2015; Bruehlheide et al., 2019).

We found a strong scale dependence of plant diversity not only across regions but also across vegetation types and biomes, as shown in Figure 3. Diversity patterns also strongly differ between the three taxonomic groups, both across biomes and across vegetation types. For vascular plants, secondary grasslands, and particularly meso-xeric grasslands, are overall the richest vegetation type, with the highest maxima in the temperate midlatitudes, where exceptionally rich meso-xeric grasslands have already been reported (Chytrý et al., 2015; Roleček et al., 2019; Hájek et al., 2020). The richest vegetation types for bryophytes and lichens strongly vary with grain size. However, it is worth reporting that communities found at high latitudes or high elevations often host the highest richness values, such as arctic-alpine heathlands, alpine grasslands and rocks and screes, following the well-documented pattern of increasing diversity of non-vascular plants towards high latitudes or elevations (Grau et al., 2007; Mateo et al., 2016; Chytrý et al., 2017). The relative importance of non-vascular plants in the total vegetation is highest in alpine grasslands and arctic-alpine heathlands and also high in sandy dry grasslands and Mediterranean grasslands. In summary, our results show that non-vascular plants can constitute a major part of the plant diversity in various habitats and thus should be more frequently considered in biodiversity studies. Moreover, we demonstrate that the richness of vascular plants is not a good proxy for fine-grain bryophyte and lichen richness, as they may behave differently, depending on vegetation type. This finding questions the use of vascular plant species richness as a surrogate for the overall diversity in open vegetation, which is suggested in several studies based on simpler assessments in forests (Pharo et al., 1999; Sætersdal et al., 2003). Likewise, Chiarucci et al. (2007) and Santi et al. (2010) found that vascular plant diversity was not a good surrogate for cryptogam diversity in any habitat type.

While the species-area relationships (SARs) were not the focus of this paper, our data illustrate some general patterns. The SARs plotted in the “semi-log” space (i.e., with area logarithmized,

but not species richness) (Figure 5 and Appendix S5, Figures S5.13-S5.16) invariably show an upward curvature, at least those that are based on the nested-plot data. This shape corresponds to a power function (see Dengler, 2008), which has recently been shown as the overall best model across the non-forest habitat types of the Palaearctic (Dengler et al., 2020b), with little variation of its exponent (z value) across grain sizes (Zhang et al., 2021). Moreover, as the example in Figure 5 illustrates, the curves only rarely cross each other, meaning that vegetation types mainly differ in their c -values (corresponding to α -diversity), while there are few systematic differences concerning z -values (corresponding to β -diversity). As shown by Dembicz et al. (2021), z -values are much more affected by disturbance regimes and heterogeneity – which are largely independent of vegetation type.

Data quality and methodological settings

GrassPlot only includes phytodiversity data that were carefully sampled with the aim of recording complete species lists within precisely delimited plots. Large vegetation-plot databases at regional and national (see Dengler et al., 2011 for an overview), continental (Chytrý et al., 2016) or global (Bruehlheide et al., 2019) scales are naturally less selective as their main aim is to get as many vegetation-plot records as possible. They include, to a large extent, traditional phytosociological relevés, in which the plot borders were often not precisely delimited in the field. Since the analysis of Chytrý (2001) of the plot data contained at that time in the Czech National Phytosociological Database, it is well-known that there are also other biases in the data. This study found, in several phytosociological classes, that the mean richness decreased above a certain threshold area, a pattern explained by the tendency of phytosociologists to select larger-than-average plots in vegetation types that are inherently poorer in species. When comparing the mean richness data from Chytrý (2001) for the three classes that are also contained in GrassPlot (*Festuco-Brometea*, *Molinio-Arrhenatheretea*, *Phragmito-Magnocaricetea*) we found substantially lower mean richness in the phytosociological database than in GrassPlot (not shown). Similarly, comparing the mean richness data of *Festuco-Brometea* grasslands from the Nordic-Baltic Grassland Vegetation Database (Dengler et al., 2006) with GrassPlot data from the same geographic region, we found a good match at 1 m², but increasing relative difference towards larger grain sizes (not shown). The consistently higher richness values in GrassPlot were

unexpected as it is often assumed that phytosociologists preferentially sample plots with a species richness above average (Holeksa & Woźniak, 2005; Diekmann et al., 2007). By contrast, most GrassPlot data are based on systematic or random sampling or the approach of the EDGG Field Workshops (Dengler et al., 2016b), which aims to maximize between-plot heterogeneity, i.e., both presumably species-rich and species-poor stands are selected for making plots (which should not bias means, but possibly increase variance). A plausible explanation for the pattern found is that the average completeness of plots in phytosociological databases is lower than most researchers, including ourselves, would have guessed. This indicates that it might be risky to take the richness data from large phytosociological databases at face value. A more comprehensive study comparing the GrassPlot benchmarks with the mean richness values derived from EVA or sPlot should explore how prevalent such a pattern is and whether its strength varies systematically between regions, vegetation types and grain sizes.

While these findings underline the good suitability of typical data contained in GrassPlot for biodiversity analyses, we do not claim that the richness records are 100% complete. It has been shown repeatedly that this is nearly impossible, even when plots are sampled by more than one experienced author (see Lepš & Hadincová, 1992; Klimeš et al., 2001; Archaux et al., 2006). However, the results support the view that the fraction of overlooked species must be minor compared to average phytosociological data and possibly even compensated by an equally minor fraction of erroneously recorded species. When the complete GrassPlot data are used, in very few cases, we also found that richness above a certain threshold appeared to stagnate or even slightly decline (Appendix S5, Figures S5.13 to S5.16). However, this can be easily explained by biases caused by large numbers of plots that were sampled in local clusters and only for one grain size but not for the others. The effect disappeared when considering only nested plot series that contain all seven standard grain sizes (Figures S5.13 to S5.16). When comparing the continuous and dashed lines in these figures, it turns out that the dashed line (the values for any plots) are largely below the continuous lines (nested plots with all the seven grain sizes). This indicates that apart from biases due to adding local clusters (which equally often should be above and below the average), even within GrassPlot data, there is a “quality gradient”: On average, the richness records in nested plots are more complete, but the differences are much

smaller than between GrassPlot and conventional phytosociological databases. Finally, also the way of recording plants as present in a plot, shoot presence vs. rooted presence (Dengler, 2008) can influence richness records as highlighted by Williamson (2003). In the habitats studied here, a visible effect occurs at grain sizes below 1 m² (Appendix S5, Table S5.2) which is consistent with findings of Güler et al. (2016), Cancellieri et al. (2017) and Zhang (2021).

While we trust that our richness data for individual plots are more reliable than most other sources, the aggregated richness patterns reported in this paper in some cases might still be biased or misleading. First, data coverage in GrassPlot is sparser than in other big vegetation-plot databases. Consequently there might be stronger biases concerning geography and vegetation types. Second, there are a few datasets in GrassPlot that have specifically been collected with the aim of studying sites of exceptional richness (e.g. Merunková et al., 2012; Roleček et al., 2014; Hájek et al., 2020). However, GrassPlot also contains data that have been sampled in regions where a certain vegetation class is known to be poorer in species than in other parts of the respective country. In addition, a prevalence of vegetation plots from one subtype of a certain category might make this entire category appear relatively richer or poorer in species than it is in reality. Similarly, not all biomes contain the same subtypes of a vegetation type. However, the magnitude of such potential biases can be assessed using the GrassPlot Diversity Explorer/Benchmarks via filtering at different levels. Third, our results are necessarily affected by the classification systems used for biomes and vegetation types. While the categories *per se* are widely used, their border might differ between different sources. For example, the relatively high mean richness values of boreal grasslands in our results are partly related to the biome classification of Schultz (2005), which includes significant areas of the forest-steppe zone in the boreal biome, while other typologies consider it a continental variant of the temperate biome (Erdős et al., 2018). Some bias may also be caused by disputed borders between vegetation types. Since the assignments to the fine-level vegetation types were largely based on syntaxonomy, and the fine-level types were fully nested in coarse categories, there are some “grey zones”, e.g. some rocky, alpine and xeric grasslands might be secondary, and, *vice versa*, some meso-xeric grasslands might be natural, particularly those in the transition to the steppic

natural grasslands (e.g., forest-steppes, Erdős et al., 2020), often maintained through grazing by wild herbivores and fire (Pärtel et al., 2015).

Potential uses and impact

Vegetation ecology. In studies on certain vegetation types, it is useful for authors to compare not only the richness values within their sample, but also to know where these are positioned in relation to the average of this vegetation type in the country or biome. Modelling studies could also benefit if they had reliable benchmarks. For example, Bruelheide et al. (2020) used a sophisticated approach to create richness maps of *Festuco-Brometea* communities in Germany at 1, 10, 100 and 1000 m² with three different approaches based on a traditional phytosociological database, but in the end, they could only “guess” which of their approaches performed best because they lacked benchmarks from more reliable data from exhaustively sampled plots.

Macroecology. An increasing number of studies use the enormous amount of vegetation-plot data from national and regional (see Dengler et al., 2011), continental (EVA; Chytrý et al., 2016) and global (sPlot; Bruelheide et al., 2019) vegetation-plot databases. This approach has great potential for macroecology as it combines fine-grain sizes with large spatial extents, a combination that could contribute to a more mechanistic understanding of patterns, but for a long time was underrepresented in macroecology (Beck et al., 2012). Moreover, vegetation-plot data allow for a much wider range of macroecological analyses than species occurrence databases do (Dengler et al., 2011; Bruelheide et al., 2019). Most of such plot-based macroecological papers take the information in the underlying databases as unquestioned facts. While such studies often address the unequal distribution of plots in space and time (Lengyel et al., 2011) and the preferential sampling of more species-rich communities (Divíšek & Chytrý, 2018), and sometimes also their different plot sizes (Večeřa et al., 2019), to our knowledge, the issue that the recorded species lists might be incomplete was hitherto not addressed in macroecological studies. Moreover, given the different traditions of phytosociology in different countries (Guarino et al., 2018), one can assume that the average degree of incompleteness might vary regionally, leading not only to biased absolute numbers but also unreliable patterns. Incomplete species lists are particularly problematic for macroecological studies on α -diversity

and some studies on β -diversity, while studies on community-weighted means of traits or assembly rules are probably less affected, at least not when assuming that the overlooked species mostly were the rare ones with low cover. Depending on the sensitivity of the study topic towards biased species richness values, macroecological users of vegetation-plot databases have several options: (i) use a subset of regions, vegetation types and/or grain sizes that, according to the comparison with the GrassPlot Diversity Benchmarks, are least affected; (ii) apply context-dependent “correction factors”; or (iii) conduct the study instead with the GrassPlot database. Depending on the question at hand, researchers should also take into account the spatial coverage and representativeness of the respective vegetation-plot databases. For example, EVA is much less spatially clustered than either GrassPlot or sPlot. In some cases, a combination of EVA and GrassPlot or sPlot and GrassPlot might be the best solution. Plots duplicated in EVA or sPlot are already documented in GrassPlot, so there is no major problem when using both sources together.

Biodiversity conservation. In conservation, a typical challenge is to prioritize areas that deserve protection. Here our benchmarks could become a useful and applicable tool. As species richness is generally seen as one of the leading criteria for such prioritizations (Brooks et al., 2006; Brum et al., 2017), one could set an objective criterion for prioritization such as plots above the third quartile or 50% above the mean value. Since the GrassPlot Diversity Benchmarks provide such values for any grain size up to 100 m² and specifically for each vegetation type, one can even compare across these categories, e.g., the threshold for alpine grasslands will be different from that for wetlands. In any case, we would like to emphasize that species richness cannot be used as a single criterion, as several naturally species-poor habitats are more species-rich after degradation, such as lower levels of salinity in saline communities. Another typical question in this context is whether a particular management or restoration measure was successful or what is the restoration potential of a specific habitat type. Did the measure achieve the typical diversity of that habitat type? Referring to richness data from the literature is troublesome in such cases as they were often recorded on different grain sizes and usually only at a single grain size, making the “translation” to another grain size challenging. All this is much easier with the GrassPlot Diversity Benchmarks, acknowledging that they largely reflect the situation during the

past two decades as there is only a small fraction of 20th-century plots included. We also acknowledge that species number should not always be used as a unique criterion for such assessments, as restoration projects often monitor richness of habitat-specific target species to avoid bias caused by sites with high richness of ruderal or alien species. Finally, we would like to advise again to carefully check plot number and spatial representativeness using the Explorer tool when using these benchmarks.

Quality check of data. In all the above-mentioned applications, the GrassPlot Diversity Explorer can be helpful for researchers and students alike to get feedback on how complete their field records likely are. The GrassPlot Diversity Benchmarks provide vegetation-plot databases with the option of checking the reliability of datasets before including them. For example, datasets with mean richness below the first quartile of the respective vegetation type × region × grain size should be considered carefully. They do not necessarily need to be excluded but could be labelled as doubtful unless the originators provide convincing reasons that the studied stands are actually so species-poor. This quality check may also be used when data from large vegetation-plot databases are selected for specific projects. Although we only provide benchmarks for eight specific grain sizes, interpolation of richness data to any other grain sizes can be easily done, as explained in Appendix S6.

Conclusions and outlook

The GrassPlot Diversity Benchmarks provide high-quality richness data from a wide range of open habitat types across the Palaearctic realm. The restriction to eight standard grain sizes, each separated by a factor of 10, is similar to some standardized sampling schemes on other continents, such as the Carolina Vegetation Survey in North America (Peet et al., 1998) and the BIOTA Observatories in Africa (Jürgens et al., 2012). Seven of the eight grain sizes are already well populated with data, only high-quality observations for 1000 m² are still sparse (which is understandable, given the enormous time effort for a complete sampling of such an area; see Dolnik, 2003). The amount of data in the underlying GrassPlot database and their spatial coverage are much lower than in the EVA (Chytrý et al., 2016) and sPlot (Bruehlheide et al., 2019) databases, which is an important constraint that may affect the aggregated patterns reflected in

the diversity benchmarks. However, we have shown that species recordings are, on average, apparently much more complete in GrassPlot. Thus, depending on the research question, either EVA/sPlot, GrassPlot or a combination of both might be the best data source. Our study further emphasizes the advantages of standardized methodologies and a set of uniform standard grain sizes.

We release this information open access in an easy-to-use format as the GrassPlot Diversity Benchmarks and in the conveniently queried online tool, the GrassPlot Diversity Explorer. Given the uneven representation of vegetation types across biomes, regions and countries, we advise users to check the number of plots available for the selected combination of region and vegetation type in order to obtain reliable benchmarks. While the GrassPlot Diversity Explorer is already quite powerful, we are considering further extensions in the future, such as an online interface to interpolate richness data between two of our standard grain sizes based on the function presented in Appendix S6, provision of the slope parameter z of species-area relationships as a β -diversity measure (see Dengler et al., 2020b), a selection option for different sampling designs and for phytosociological units below the class level, or maps of mean and maximum richness. While the provided open access data will suffice for many purposes, it is also possible to approach the GrassPlot Consortium with a project proposal (according to the GrassPlot Bylaws; available at <https://edgg.org/databases/GrassPlot>) for individual plot-based richness records together with environmental data. Last but not least, GrassPlot is a “living database”, which is constantly enriched and improved. Those who would like to join the GrassPlot Consortium with suitable data can approach the first author (IB), who is serving as a database manager. The GrassPlot Diversity Explorer will be dynamically updated with the newest versions of GrassPlot, while a new fixed version of the GrassPlot Diversity Benchmarks is planned to be released regularly.

Acknowledgements

We thank Manuel J. Steinbauer for the concept of the richness map in Figure 2. We thank the hundreds of vegetation ecologists who sampled the high-quality data used in this article and contributed them to GrassPlot.

Author contributions

JD and IB conceived the research idea; IB compiled the data and managed the database; IB and JD wrote the paper; RPi developed the GrassPlot Diversity Explorer and ID the GrassPlot Diversity Benchmarks; ID, FG, LK, CorM, TR and KVM created the figures; RG managed the author list; IAx, SBu, SBa, SBo, HHB, MCh, TCo, PDF, FE, MHá, BJA, AKu, ZM, RJP, MP, RPä, RPi, ZP, HCP, JR, LS, MTe, KVM and JWu discussed the results and commented on the manuscript; all authors contributed data and approved the manuscript.

Data availability statement

The aggregated data (as used in this paper) for any combination of the taxonomic group, grain size, vegetation type, region, biome, and methodological settings (rooted vs. shoot; subsetting to only those plots belonging to nested series with at least seven standard grain sizes) are provided in Appendix S3. Future updates will be made available as GrassPlot Diversity Benchmarks (fixed versions) and dynamically in the GrassPlot Diversity Explorer (both at <https://edgg.org/databases/GrasslandDiversityExplorer>). The underlying plot-level data are available upon request from the GrassPlot database, following its Bylaws (<https://edgg.org/databases/GrassPlot>).

References

- Allaire, J. (2019) *rconnect: Deployment interface for R markdown documents and Shiny applications*. R package version 0.8.16.
- Archaux, F., Gosselin, F., Bergès, L. & Chevalier, R. (2006) Effects of sampling time, quadrat richness and observer on exhaustiveness of plant censuses. *Journal of Vegetation Science*, 17, 299–306. <https://doi.org/10.1111/j.1654-1103.2006.tb02449.x>
- Axmanová, I., Kalusová, V., Danihelka, J., Dengler, J., Pergl, J., Pyšek, P. et al. (2021) Neophyte invasions in European grasslands. *Journal of Vegetation Science*, 32, e12994.
- Barthlott, W., Mutke, J., Rafiqpoor, D., Kier, G. & Kreft, H. (2005) Global centers of vascular plant diversity. *Nova Acta Leopoldina NF*, 92(342), 61–83. <https://doi.org/10.1111/jvs.12994>

Barthlott, W., Hostert, A., Kier, G., Küpfer, W., Kreft, H., Mutke, J. et al. (2007) Geographic patterns of vascular plant diversity at continental to global scales. *Erdkunde*, 61, 305–315. <https://doi.org/10.3112/erdkunde.2007.04.01>

Beck, J., Ballesteros-Mejia, L., Buchmann, C.M., Dengler, J., Fritz, S.A., Gruber, B. et al. (2012) What's on the horizon for macroecology? *Ecography*, 35, 673–683. <https://doi.org/10.1111/j.1600-0587.2012.07364.x>

Biurrun, I., Burrascano, S., Dembicz, I., Guarino, R., Kapfer, J., Pielech, R. et al. (2019) GrassPlot v. 2.00 – first update on the data-base of multi-scale plant diversity in Palaeartic grasslands. *Palaeartic Grasslands*, 44, 26–47. <https://doi.org/10.21570/EDGG.PG.44.26-47>

Bostock, M. Rodden, K., Warne, K., Russell, K, Breitwieser, F. & Yetman, C.J. (2020) *sunburstR: Sunburst 'Htmlwidget'*. R package version 2.1.5.

Brooks, T. M., Mittermeier, R. A., da Fonseca, G. A. B., Gerlach, J., Hoffman, M., Lamoreux, J. F. et al. (2006) Global biodiversity conservation priorities. *Science*, 313, 58–61. <https://doi.org/10.1126/science.1127609>

Bruelheide, H., Dengler, J., Purschke, O., Lenoir, J., Jiménez-Alfaro, B., Hennekens, S.M. et al. (2018) Global trait–environment relationships of plant communities. *Nature Ecology & Evolution*, 2, 1906–1917. <https://doi.org/10.1038/s41559-018-0699-8>

Bruelheide, H., Dengler, J., Jiménez-Alfaro, B., Purschke, O., Hennekens, S.M., Chytrý, M. et al. (2019) sPlot – A new tool for global vegetation analyses. *Journal of Vegetation Science*, 30, 161–186. <https://doi.org/10.1111/jvs.12710>

Bruelheide, H., Jiménez-Alfaro, B., Jandt, U. & Sabatini, F.M. (2020) Deriving site-specific species pools from large databases. *Ecography*, 43, 1–14. <https://doi.org/10.1111/ecog.05172>

Brum, F. T., Graham, C. H., Costa, G. C., Hedges, S. B., Penone, C., Radeloff, V. C. et al. (2017) Global priorities for conservation across multiple dimensions of mammalian diversity. *Proceedings of the National Academy of Sciences of the United States of America*, 114, 7641–7646. <https://doi.org/10.1073/pnas.1706461114>

Brummitt, N., Araújo, A.C. & Harris, T. (2020) Areas of plant diversity—What do we know? *Plants People Planet*, 3, 33–44. <https://doi.org/10.1002/ppp3.10110>

Cancellieri, L., Mancini, L. D., Sperandii, M. G. & Filibeck, G. (2017) In and out: effects of shoot- vs. rooted-presence sampling methods on plant diversity measures in mountain grasslands. *Ecological Indicators*, 72, 315–321. <https://doi.org/10.1016/j.ecolind.2016.08.029>

Chang, W., Cheng, J., Allaire, J., Xie, Y. & McPherson, J. (2020) *shiny: Web application framework for R*. R package version 1.5.0.

Cheng, J., Karambelkar, B. & Xie, Y. (2019) *leaflet: Create interactive web maps with the JavaScript 'Leaflet' library*. R package version 2.0.3.

Chiarucci, A., D'Auria F. & Bonini, I. (2006) Is vascular plant species diversity a predictor of bryophyte species diversity in Mediterranean forests? *Biodiversity and Conservation*, 16, 525–545.

Chytrý, M. (2001) Phytosociological data give biased estimates of species richness. *Journal of Vegetation Science*, 12, 459–444. <https://doi.org/10.1111/j.1654-1103.2001.tb00190.x>

Chytrý, M. & Otýpková, Z. (2003) Plot sizes used for phytosociological sampling of European vegetation. *Journal of Vegetation Science*, 14, 563–570. <https://doi.org/10.1111/j.1654-1103.2003.tb02183.x>

Chytrý, M., Dražil, T., Hájek, M., Kalníková, V., Preislerová, Z., Šibík, J. et al. (2015) The most species-rich plant communities in the Czech Republic and Slovakia (with new world records). *Preslia*, 97, 217–278.

Chytrý, M., Hennekens, S.M., Jiménez-Alfaro, B., Knollová, I., Dengler, J., Jansen, F. et al. (2016) European Vegetation Archive (EVA): An integrated database of European vegetation plots. *Applied Vegetation Science*, 19, 173–180. <https://doi.org/10.1111/avsc.12191>

Chytrý, M., Horsák, M., Syrovátka, V., Danihelka, J., Ermakov, N., German, D. A. et al. (2017) Refugial ecosystems in central Asia as indicators of biodiversity change during the Pleistocene–Holocene transition. *Ecological Indicators*, 77, 357–367. <https://doi.org/10.1016/j.ecolind.2016.12.033>

Comtois, D. (2020) *summarytools: Tools to quickly and neatly summarize data*. R package version 0.9.6.

Dembicz, I., Dengler, J., Steinbauer, M. J., Matthews, T. J., Bartha, S., Burrascano, S. et al. (2021) Fine-grain beta diversity of Palaeartic grassland vegetation. *Journal of Vegetation Science*, 32, e13045. <https://doi.org/10.1111/jvs.13045>

Dengler, J. (2008) Pitfalls in small-scale species-area sampling and analysis. *Folia Geobotanica*, 43, 269–287. <https://doi.org/10.1007/s12224-008-9014-9>

Dengler, J., Rusina, S., Boch, S., Bruun, H. H., Diekmann, M., Dierßen, K. et al. (2006) Working group on dry grasslands in the Nordic and Baltic region – outline of the project and first results for the class *Festuco-Brometea*. *Annali di Botanica, Nuova Serie*, 6, 1–28.

Dengler, J., Jansen, F., Glöckler, F., Peet, R.K., De Cáceres, M., Chytrý, M. et al. (2011) The Global Index of Vegetation-Plot Databases (GIVD): a new resource for vegetation science. *Journal of Vegetation Science*, 22, 582–597. <https://doi.org/10.1111/j.1654-1103.2011.01265.x>

Dengler, J., Biurrun, I., Apostolova, I., Baumann, E., Becker, T., Berastegi, A. et al. (2016a) Scale-dependent plant diversity in Palaeartic grasslands: a comparative overview. *Bulletin of the Eurasian Dry Grassland Group*, 31, 12–26.

Dengler, J., Boch, S., Filibeck, G., Chiarucci, A., Dembicz, I., Guarino, R. et al. (2016b) Assessing plant diversity and composition in grasslands across spatial scales: The standardised EDGG sampling methodology. *Bulletin of the Eurasian Grassland Group*, 32, 13–30.

Dengler, J., Wagner, V., Dembicz, I., García-Mijangos, I., Naqinezhad, A., Boch, S. et al. (2018) GrassPlot – a database of multi-scale plant diversity in Palaeartic grasslands. *Phytocoenologia*, 48, 331–347. <https://doi.org/10.1127/phyto/2018/0267>

Dengler, J., Biurrun, I., Boch, S., Dembicz, I. & Török, P. (2020a) Grasslands of the Palaeartic biogeographic realm: introduction and synthesis. In: Goldstein, M.I. and DellaSala, D.A. (Eds.), *Encyclopedia of the world's biomes*. Amsterdam: Elsevier, pp. 617–637.

- Dengler, J., Matthews, T. J., Steinbauer, M. J., Wolfrum, S., Boch, S., Chiarucci, A. et al. (2020b) Species–area relationships in continuous vegetation: Evidence from Palaeartic grasslands. *Journal of Biogeography*, 47, 72–86. <https://doi.org/10.1111/jbi.13697>
- Diekmann, M., Kühne, A. & Isermann, M. (2007) Random vs non-random sampling: Effects on patterns of species abundance, species richness and vegetation–environment relationships. *Folia Geobotanica*, 42, 179–190. <https://doi.org/10.1007/BF02893884>
- Divíšek, J. & Chytrý, M. (2018) High-resolution and large-extent mapping of plant species richness using vegetation-plot databases. *Ecological Indicators*, 89, 840–851. <https://doi.org/10.1016/j.ecolind.2017.11.005>
- Dolnik, C. (2003) Species-area relations of forest and open habitats – A contribution to the recording of plant biodiversity with special attention to lichens and mosses using the example of the National Park Kurische Nehrung (Russia) (German). *Mitteilungen der Arbeitsgemeinschaft Geobotanik in Schleswig-Holstein und Hamburg*, 62, 1–183.
- Erdős, L., Ambarlı, D., Anenkhonov, O. A., Bátori, Z., Cserhalmi, D., Kiss, M. et al. (2018) The edge of two worlds: A new review and synthesis on Eurasian forest-steppes. *Applied Vegetation Science*, 21, 345–362. <https://doi.org/10.1111/avsc.12382>
- Erdős, L., Török, P., Szitár, K., Bátori, Z., Tölgyesi, C., Kiss, P. J. et al. (2020). Beyond the forest-grassland dichotomy: The gradient-like organization of habitats in forest-steppes. *Frontiers in Plant Science*, 11, 236. <https://doi.org/10.3389/fpls.2020.00236>
- Faber-Langendoen, D. & Josse, C. (2010) *World grasslands and biodiversity patterns*. Arlington, VA: NatureServe.
- Fischer, M., Rounsevell, M., Torre-Marín Rando, A., Mader, A., Church, A., Elbakidze, M. et al. (eds.) (2018) *Summary for policymakers of the regional assessment report on biodiversity and ecosystem services for Europe and Central Asia of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services*. Bonn: IPBES secretariat.
- Gaston, K. J. (2000) Global patterns in biodiversity. *Nature*, 405, 220–227.

Geffert, J. L., Frahm, J.-P., Barthlott, W. & Mutke, J. (2013) Global moss diversity: spatial and taxonomic patterns of species richness. *Journal of Bryology*, 35, 1–11. <https://doi.org/10.1179/1743282012Y.0000000038>

Grau, O., Grytnes, J. A. & Birks, H. J. B. (2007) A comparison of altitudinal species richness patterns of bryophytes with other plant groups in Nepal, Central Himalaya. *Journal of Biogeography*, 34, 1907–1915. <https://doi.org/10.1111/j.1365-2699.2007.01745.x>

Guarino, R., Willner, W., Pignatti, S., Attorre, F. & Loidi, J. (2018) Spatio-temporal variations in the application of the Braun-Blanquet approach in Europe. *Phytocoenologia*, 48, 239–250. <https://doi.org/10.1127/phyto/2017/0181>

Güler, B., Jentsch, A., Bartha, S., Bloor, J. M. G., Campetella, G., Canullo, R. et al. (2016) How plot shape and dispersion affect plant species richness counts: implications for sampling design and rarefaction analyses. *Journal of Vegetation Science*, 27, 692–703. <https://doi.org/10.1111/jvs.12411>

Hájek, M., Hájková, P. & Roleček, J. (2020) A novel dataset of permanent plots in extremely species-rich temperate grasslands. *Folia Geobotanica*, 55, 257–268. <https://doi.org/10.1007/s12224-020-09372-6>

Holeksa, J. & Woźniak, G. (2005) Biased vegetation patterns and detection of vegetation changes using phytosociological databases. A case study in the forests of the Babia Góra National Park (the West Carpathians, Poland). *Phytocoenologia*, 35, 1–18.

Hooper, D. U., Chapin III, F. S., Ewel, J. J., Hector, A., Inchausti, P., Lavorel, S. et al. (2005) Effects of biodiversity on ecosystem functioning: A consensus of current knowledge. *Ecological Monographs*, 75, 3–35. <https://doi.org/10.1890/04-0922>

Jürgens, N., Schmiedel, U., Haarmeyer, D.H., Dengler, J., Finckh, M., Goetze, D. et al. (2012) The BIOTA Biodiversity Observatories in Africa – A standardized framework for large-scale environmental monitoring. *Environmental Monitoring and Assessment*, 184, 655–678. <https://doi.org/10.1007/s10661-011-1993-y>

Kassambara, A. (2020) *ggpubr: 'ggplot2' Based Publication Ready Plots*. R package version 0.4.0.

Kier, G., Mutke, J., Dinerstein, E., Ricketts, T. H., Küper, W., Kreft, H. & Barthlott, W. (2005) Global patterns of plant diversity and floristic knowledge. *Journal of Biogeography*, 32, 1107–1116. <https://doi.org/10.1111/j.1365-2699.2005.01272.x>

Klímeš, L., Dančák, M., Hájek, M., Jongepierová, I. & Kučera, T. (2001) Scale-dependent biases in species counts in a grassland. *Journal of Vegetation Science*, 12, 699–704. <https://doi.org/10.2307/3236910>

Körner, C., Jetz, W., Paulsen, J., Payne, D., Rudmann-Maurer, K. & Spehn, E. M. (2017) A global inventory of mountains for bio-geographical applications. *Alpine Botany*, 127, 1–15. <https://doi.org/10.1007/s00035-016-0182-6>

Küper, W., Sommer, J. H., Lovett, J. C., Mutke, J., Linder, H. P., Beentje, H. J. et al. (2005) Africa's hotspots of biodiversity redefined. *Annals of the Missouri Botanical Garden*, 91, 525–535.

Lengyel, A., Chytrý, M. & Tichý, L. (2011) Heterogeneity-constrained random resampling of phytosociological databases. *Journal of Vegetation Science*, 22, 175–183. <https://doi.org/10.1111/j.1654-1103.2010.01225.x>

Lepš, J. & Hadincová, V. (1992) How reliable are our vegetation analyses? *Journal of Vegetation Science*, 3, 119–124. <https://doi.org/10.2307/3236006>

Mateo, R. G., Broennimann, O., Normand, S., Petitpierre, B., Araújo, M.B., Svenning, J.C. et al. (2016) The mossy north: an inverse latitudinal diversity gradient in European bryophytes. *Scientific Reports*, 6, e25546. <https://doi.org/10.1038/srep25546>

McKerrow, A. J., Tarr, N. M., Rubino, M. J. and Williams, S. G. (2018) Patterns of species richness hotspots and estimates of their protection are sensitive to spatial resolution. *Diversity and Distributions*, 24, 1464–1377. <https://doi.org/10.1111/ddi.12779>

Merunková, K., Preislerová, Z. & Chytrý, M. (2012) White Carpathian grasslands: can local ecological factors explain their extraordinary species richness? *Preslia*, 84, 311–325.

Myers, N., Mittermeier, R. A., Mittermeier, C. G., da Fonseca, G. A. B. & Kent, J. (2000) Biodiversity hotspots for conservation priorities. *Nature*, 403, 853–858.

Pärtel, M., Bruun, H. H. & Sammul, M. (2005) Biodiversity in temperate European grasslands: origin and conservation. *Grassland Science in Europe* 10, pp. 1–14.

Pärtel, M., Bennett, J. A. & Zobel, M. (2016) Macroecology of biodiversity: disentangling local and regional effects. *New Phytologist*, 211, 404–410. <https://doi.org/10.1111/nph.13943>

Peet, R.K., Wentworth, T.R. & White, P.S. (1998) A flexible, multipurpose method for recording vegetation composition and structure. *Castanea*, 63, 262–274.

Perrier, V., Meyer, F. & Granjon, D. (2020) *shinyWidgets: custom inputs widgets for shiny*. R package version 0.5.3.

Pharo, E. J., Beattie, A. J. & Binns, D. (1999) Vascular plant diversity as a surrogate for bryophyte and lichen diversity. *Conservation Biology*, 13, 282–292. <https://doi.org/10.1046/j.1523-1739.1999.013002282.x>

R Core Team (2020) *R: A language and environment for statistical computing*. Vienna: R Foundation for Statistical Computing.

Rahbek, C. (2005) The role of spatial scale and the perception of large-scale species-richness patterns. *Ecology Letters*, 8, 224–239. <https://doi.org/10.1111/j.1461-0248.2004.00701.x>

Reid, W. V. (1998) Biodiversity hotspots. *Trends in Ecology & Evolution*, 13, 275–280. [https://doi.org/10.1016/S0169-5347\(98\)01363-9](https://doi.org/10.1016/S0169-5347(98)01363-9)

Roleček, J., Čornej, I.I. & Tokarjuk, A.I. (2014) Understanding the extreme species richness of semi-dry grasslands in east-central Europe: a comparative approach. *Preslia*, 86, 5–27.

Roleček, J., Dřevojan, P., Hájková, P. & Hájek, M. (2019) Report of new maxima of fine-scale vascular plant species richness recorded in East-Central European semi-dry grasslands. *Tuexenia*, 39, 423–431. <https://doi.org/10.14471/2019.39.008>

Sali, A. & Attali, D. (2020) *shinycssloaders: add loading animations to a 'shiny' output while it's recalculating*. R package version 1.0.0.

- Santi, E., Maccherini, S., Rocchini, D., Bonini, I., Brunialti, G., Favilli, L. et al. (2010) Simple to sample: Vascular plants as surrogate group in a nature reserve. *Journal for Nature Conservation*, 18, 2–11. <https://doi.org/10.1016/j.jnc.2009.02.003>
- Sætersdal, M., Gjerde, I., Blom, H. H., Ihlen, P. G., Myrseth, E. W., Pommeresche, R. et al. (2003) Vascular plants as a surrogate species group in complementary site selection for bryophytes, macrolichens, spiders, carabids, staphylinids, snails, and wood living polypore fungi in a northern forest. *Biological Conservation*, 115, 21–31. [https://doi.org/10.1016/S0006-3207\(03\)00090-9](https://doi.org/10.1016/S0006-3207(03)00090-9)
- Schultz, J. (2005) *The ecozones of the world. The ecological division of the geosphere*. 2nd edition. Berlin: Springer.
- Shriner, S. A., Wilson, K. R. & Flather, C. H. (2006) Reserve networks based on richness hotspots and representation vary with scale. *Ecological Applications*, 16, 1660–1673. [https://doi.org/10.1890/1051-0761\(2006\)016\[1660:RNBORH\]2.0.CO;2](https://doi.org/10.1890/1051-0761(2006)016[1660:RNBORH]2.0.CO;2)
- Tilman, D. & Downing, J. A. (1994) Biodiversity and stability in grasslands. *Nature*, 367, 363–365. <https://doi.org/10.1038/367363a0>
- Večeřa, M., Divíšek, J., Lenoir, J., Jiménez-Alfaro, B., Biurrun, I., Knollová, I. et al. (2019) Alpha diversity of vascular plants in European forests. *Journal of Biogeography*, 46, 1919–1935. <https://doi.org/10.1111/jbi.13624>
- Whittaker, R. J., Araújo, M. B., Jepson, P., Ladle, R. J., Watson, J. E. M. & Willis, K. J. (2015) Conservation Biogeography: assessment and prospect. *Diversity and Distributions*, 11, 3–23. <https://doi.org/10.1111/j.1366-9516.2005.00143.x>
- Wickham, H. (2016) *ggplot2: Elegant Graphics for Data Analysis*. 2nd edition. New York: Springer.
- Wickham, H. & Henry, L. (2020) *tidyr: Tidy Messy Data*. R package version 1.1.0.
- Wickham, H., François, R., Henry, L. & Müller, K. (2020) *dplyr: A Grammar of Data Manipulation*. R package version 1.0.0.
- Williamson, M. (2003) Species-area relationships at small scales in continuum vegetation. *Journal of Ecology*, 91, 904–907. <https://doi.org/10.1046/j.1365-2745.2003.00816.x>

Wilson, J. B., Peet, R. K., Dengler, J. & Pärtel, M. (2012) Plant species richness: the world records. *Journal of Vegetation Science*, 23, 796–802. <https://doi.org/10.1111/j.1654-1103.2012.01400.x>

Yen, J. D. L., Dorrough, J., Oliver, I., Somerville, M., McNellie, M. J., Watson, C. J. & Vesk, P. A. (2019) Modeling biodiversity benchmarks in variable environments. *Ecological Applications*, 29, e01970. <https://doi.org/10.1002/eap.1970>

Zhang, J., Gillet, F., Bartha, S., Alatalo, J.M., Biurrun, I., Dembicz, I. et al. (2021) Scale dependence of species-area relationships is widespread but generally weak in Palaeartic grasslands. *Journal of Vegetation Science*, 32, e13044. <https://doi.org/10.1111/jvs.13044>

Accepted Article

Supporting Information

Additional supporting information may be found online in the Supporting Information section.

Appendix S1. Overview of the 225 datasets included in GrassPlot v2.10 and their references.

Appendix S2. Complementary information on the data.

Appendix S3. The GrassPlot Diversity Benchmarks, version 2.10.

Appendix S4. Additional richness maps.

Appendix S5. Additional detailed results.

Appendix S6. Interpolation of richness data to other grain sizes.

Appendix S7. Funding information.

Accepted Article

Tables

Table 1. The number of available plots per taxonomic group and grain sizes. Standard sizes are indicated; 0.001 m² also includes 0.0009 m²; 0.1 m² includes 0.09 m²; 10 m² includes 9 and 10.89 m²; and 1000 m² includes 900 and 1024 m². N_{all} = total number of plots. $N_{\text{ind.}}$ = number of independent observations, i.e., after averaging several subplots of the same grain size in the same nested series.

Grain size [m ²]	All groups		Vascular plants		Bryophytes		Lichens	
	N_{all}	$N_{\text{ind.}}$	N_{all}	$N_{\text{ind.}}$	N_{all}	$N_{\text{ind.}}$	N_{all}	$N_{\text{ind.}}$
0.0001	1,959	774	2,904	1,440	1,973	781	1,959	774
0.001	1,900	707	4,341	1,727	1,914	714	1,900	707
0.01	2,653	877	70,548	4,875	2,688	905	6,820	1,876
0.1	1,925	717	5,756	3,788	1,939	724	1,925	717
1	2,821	1,583	25,432	20,500	3,044	1,799	3,002	1,764
10	2,924	2,171	11,035	9,658	4,599	3,839	2,925	2,172
100	1,336	1,316	6,321	6,274	1,363	1,343	1,436	1,416
1000	45	45	187	187	45	45	45	45
All sizes	15,563	8,190	126,524	48,449	17,565	10,150	20,012	9,471

Table 2. Maximum richness values for each taxonomic group and grain size across coarse-level vegetation types. The highest values for each taxonomic group are shown in bold. A: natural grasslands; B: secondary grasslands; C: azonal communities; D: dwarf shrublands; E: tall-forb and ruderal communities; F: deserts and semi-deserts. + or - before the maximum values indicate that they are derived from slightly smaller (+) or bigger (-) grain sizes than the standard ones, i.e., 0.0009, 0.09, 9, 10.89, 900 or 1024 m², respectively. Maximum richness for the exact grain size, if available, is indicated in brackets in the upper case. No filtering by sampling method (rooted vs. shoot) was applied.

Area [m ²]	Complete vegetation						Vascular plants					
	A	B	C	D	E	F	A	B	C	D	E	F
0.0001	10	10	4	6	7	1	8	11	4	5	7	4
0.001	14	19	8	7	10	3	12	19	7	7	9	5
0.01	25	29	11	10	18	7	21	24	10	18	16	10
0.1	43	46	20	35	30	12	34	43	16	34	28	18
1	63	82	39	39	49	20	59	79	33	49	44	29
10	90	101	64	69	71	51	86	-106⁽⁹⁸⁾	68	65	68	+48⁽⁴⁷⁾
100	128	159	65	93	123	48	119	144	82	113	122	71
1000	-	+123⁽⁵⁸⁾	+89	-	-	-	134	+97⁽⁸³⁾	+82⁽³⁰⁾	96	126	95

Area [m ²]	Bryophytes						Lichens					
	A	B	C	D	E	F	A	B	C	D	E	F
0.0001	5	5	3	2	2	1	2	4	1	0	0	1
0.001	+9⁽⁶⁾	8	3	2	3	+1 ⁽⁰⁾	+6⁽²⁾	+5⁽³⁾	1	1	0	+1 ⁽⁰⁾
0.01	18	10	3	3	4	1	8	7	2	3	0	1
0.1	+24⁽⁹⁾	+14⁽¹⁰⁾	4	4	8	+1 ⁽⁰⁾	+15⁽⁸⁾	+10⁽⁵⁾	4	2	2	+1 ⁽⁰⁾
1	31	18	10	11	8	2	21	17	6	13	3	2
10	+40⁽¹⁸⁾	+22⁽¹⁹⁾	27	10	16	+11 ⁽¹⁾	+24⁽²³⁾	+20⁽¹²⁾	7	2	3	+10 ⁽¹⁾
100	38	32	19	16	21	1	25	31	15	13	4	0

Accepted Article

Figures




Figure 1. Distribution of plots in the Palearctic realm. Biomes are shown in different colours, pie-charts show the fraction of vegetation types represented by the plots (black dots) included in each biome. The category “Others” includes vegetation types represented by less than 2% plots in each particular biome. C.4 Saline steppes and semi-deserts and F.1 Alpine deserts do not reach this threshold in any biome. The box plots show the elevation distribution of plots across biomes, with the number of plots (n) above each bar. To fill in the Arabian Peninsula, the biome Tropics with summer rain is indicated in orange colour although GrassPlot does not contain any data from this biome.




Figure 2. Richness hotspots and coldspots of vascular plants across spatial grains in grasslands and other open habitats across the Palearctic realm. Concentric circles represent mean species richness at grain sizes 0.01, 1 and 100 m² within hexagons of ca. 5° x 5°, from the lowest (blue) to the highest (red), while white indicates the absence of data of that grain size.




Figure 3. Richness of vascular plants across coarse-level vegetation types and biomes for the four most represented grain sizes (0.1 m², 1 m², 10 m², 100 m²). Under each bar, the number of plots is given. No filtering by sampling method (rooted vs. shoot) was applied.




Figure 4. Richness at 1 m² of the four taxonomic groups across coarse-level vegetation types. Under each bar, the number of plots is given. No filtering by sampling method (rooted vs. shoot) was applied.

Vegetation type

- A.1 Alpine grasslands
- A.4 Rocky grasslands
- B.2 Meso-xeric grasslands
- ▲ A.3 Xeric grasslands and steppes
- ◆ B.1 Sandy dry grasslands
- * B.5 Mediterranean grasslands




Figure 5. Species-area relationships for vascular plants (a) and complete vegetation (b) for six selected grassland types. Only plots belonging to nested series with at least seven standard grain sizes were included. No filtering by sampling method (rooted vs. shoot) was applied.




Figure 6. Screenshots presenting selected functionalities of the GrassPlot Diversity Explorer. Top: SARs for six selected vegetation types in the panel Species-area relationships; bottom: map showing the overall plot distribution in Western, Central and Southern Europe in the panel Maps.