

RESEARCH

Open Access



The impact of periodontal disease on the clinical outcomes of COVID-19: A systematic review and meta-analysis

Sadeq Ali Al-Maweri^{1*}, Mohammed Nasser Alhajj^{2,3}, Esam Halboub⁴, Faleh Tamimi¹, Nosizana Mohd Salleh², Mohammed Sultan Al-Ak'hali⁵, Saba Kassim⁶, Saleem Abdulrab⁷, Lamyia Anweigi¹ and Marwan Mansoor Ali Mohammed⁸

Abstract

Background A possible relationship between periodontitis (PD) and COVID-19 and its adverse outcomes has been suggested. Hence, the present systematic review and meta-analysis aimed to investigate the available evidence regarding the potential association between periodontitis (PD) and COVID-19 and its adverse outcomes.

Materials and methods PubMed, Scopus, Web of Science, and Google Scholar were searched for relevant studies published up to April 15th, 2023. Studies that evaluated the association between PD and COVID-19 were included. Risk of bias was evaluated by two reviewers, and meta-analyses were performed using RevMan 5.3 software.

Results A total of 22 studies involving 92,535 patients from USA, Europe, Asia, the Middle East and South America were included; of these, 12 were pooled into the meta-analysis. Most of the studies (19 studies) reported a significant association between PD and COVID-19. The pooled data found a significant association between PD and COVID-19 outcomes: more severe symptoms (OR = 6.95, $P = 0.0008$), ICU admissions (OR = 3.15, $P = 0.0001$), and mortality (OR = 1.92, $P = 0.21$). Additionally, compared to mild PD, severe PD was significantly associated with higher risks of severe COVID-19 outcomes: severe symptoms ($P = 0.02$); ICU admission ($P = 0.0001$); and higher mortality rates ($P = 0.0001$). The results also revealed 58% higher risk for COVID-19 infection in patients with PD ($P = 0.00001$).

Conclusions The present findings suggest a possible association between poor periodontal health and the risk of poor COVID-19 outcomes. However, owing to the observed methodological heterogeneity across the included studies, further prospective cohort studies with standardized methodologies are warranted to further unravel the potential association between periodontal disease and COVID-19 and its adverse outcomes.

Keywords COVID-19 outcomes, Periodontal disease, Association, Systematic review

*Correspondence:

Sadeq Ali Al-Maweri
Sadali05@hotmail.com

Full list of author information is available at the end of the article



© The Author(s) 2023. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

Background

The Corona virus disease of 2019 (COVID-19) caused by the SARS-CoV-2 virus has resulted in an enormous impact on the global health and economy [1–3]. Despite the fact that most COVID-19 patients recover without major complications, few patients still suffer severe complications including multiple organ failure and death [4, 5]. Such complications are driven by serious conditions triggered by the infection such severe acute respiratory distress, systemic inflammatory reactions, and coagulopathies [6, 7]. It became obvious that several comorbidities such as obesity, hypertension, diabetes and advanced age are associated with severe COVID-19 [4–6, 8]. A considerable fraction of apparently healthy and young patients, with no clear identifiable risk factors, however, still develops severe complications.

Periodontitis (PD), the most common form of periodontal diseases, is a chronic disease involving the inflammation and subsequent damage of the tissues surrounding the teeth [9]. If not treated properly, PD leads to the destruction of the bone surrounding the teeth and ultimately loss of the teeth themselves [10]. Beyond such local consequences, PD can also have detrimental effects on systemic health [10]. Growing evidence links PD to several systemic diseases including diabetes mellitus, cardiovascular diseases, pneumonia, metabolic syndrome, and cancers [9, 11–17]. Such systemic effects of PD might be ascribed to the bacterial products and inflammatory mediators in the periodontal infected tissues that can reach the blood stream and increase the systemic inflammatory burden [18]. Worth to mention that periodontal pathogens can reach the respiratory tract, aggravating, and/or even initiating respiratory infections [17].

Given that PD shares with COVID-19 many features including comorbidities and increased blood levels of inflammation and coagulation biomarkers, [19, 20] several researchers have hypothesized that PD could be associated with a higher risk of COVID-19, and the development of its adverse outcomes [20, 21]. Marouf et al. reported a higher risk of COVID-19 complications including death (OR=8.81), intensive care unit (ICU) admission (OR=3.54), and the need for assisted ventilation (OR=4.57) in patients with periodontitis [22]. Gupta et al. reported that the likelihood of requiring assisted ventilation, hospital admission, death, and getting COVID-19-related pneumonia were 7.45, 36.52, 14.58, and 4.42 folds, respectively, in COVID-19 patients with PD compared to COVID-19 patients without PD [23]. Additionally, 2022 case–control study among US COVID-19 patients reported significantly greater missing teeth and alveolar loss tooth in Covid-19 positive patients and in those with severe complications [24]. Interestingly, a 2022 case–control study by Said et al. reported

a significant association between history of periodontal therapy and favorable COVID-19 outcomes, indicating a positive role of periodontal therapy on Covid-19 complications [25]. Using a two-sample Mendelian randomization, Wang et al. revealed that PD was significantly associated with higher risk for COVID-19 infection and severe complications [26]. Conversely, two studies with a large sample size from UK Biobank cohorts did not support any significant association between PD and the risk and outcomes of COVID-19 [27, 28]. Similarly, a retrospective cross-sectional Dutch study did not find a significant association between alveolar bone loss and complications of COVID-19 [29]. One systematic review of two studies reported a significant association between PD and COVID-19 [30]. A more recent systematic review and meta-analysis by Baima et al., [31] investigating the potential link between PD and Covid-19, reported a significant association between PD and Covid-19 susceptibility and poor outcomes [31]. Nevertheless, the latter review included only very limited studies (n=8; studies published up till Feb. 2022), and since then numerous relevant studies were published with interesting findings. Additionally, the latter two systematic reviews did not in-depth assess the correlation between severity of PD and severity of Covid-19 symptoms. Therefore, this systematic review and meta-analysis sought to comprehensively analyze and summarize the available evidence on the association between PD and COVID-19. The focused question was: Does periodontal health status have an impact on COVID-19 risk and clinical outcomes?

Materials and methods

The present systematic review and meta-analysis adhered strictly to the PRISMA 2020 guidelines and followed utterly PECO principles. This systematic review was registered in Open Science Framework (OSF) registries (<https://doi.org/10.17605/OSF.IO/KW7TC>). The PECO research question was: Is periodontal disease a risk factor for COVID-19 and its poor outcomes?

Eligibility criteria

All studies (cohort, case–control, and cross-sectional and interventional studies) that assessed the association of periodontal diseases with COVID-19 outcomes in humans were eligible.

Exposure: periodontal disease parameters.

Outcome: COVID-19 infection and/or its associated adverse outcomes.

Case reports, post-mortem studies, animal and experimental studies, review articles, commentaries, and studies with unclear exposures/outcomes were excluded.

Search strategy and information sources

An extensive search of online databases (PubMed, Scopus, and Web of Science) and Google Scholar search engine, supplemented with manual search was conducted independently by two reviewers (SA and MA) on April 16th, 2023 for all relevant studies published from December 2019 up to April 15th, 2023. The following Mesh terms and free keywords were used: ("Periodontal Diseases"[Mesh] OR "Oral Health"[Mesh] OR "Periodontal disease" OR periodontitis" OR Periodont* OR periodontal pathogen*) AND ("SARS-CoV-2"[Mesh] OR COVID-19). Detailed information about the search strategies is presented in Supplementary Table 1.

Screening and selection process

All retrieved studies were exported to the EndNote program, which eased removal of duplicates. Then, the titles and abstracts of all articles were screened independently by two reviewers (SA and FT), and irrelevant articles were excluded. The full-texts of all potentially eligible articles were independently evaluated by the two reviewers for inclusions, and irrelevant articles were eliminated.

Data extraction

All relevant data were independently extracted by two investigators (SA, MA) using customized forms. The extracted data included the following: first author name, country of the study, study design, sample size, age and gender of the participants, periodontal variables (exposure) and COVID-19 variables (the main outcomes).

Quality assessment

Two reviewers (SA and MA) assessed independently the quality of all included studies using the Newcastle Ottawa Scale (NOS) for assessing the quality of non-randomized studies [32]. Disagreements, if any, were resolved by discussion. The overall quality of each study was rated as either: high quality, 7–9 stars; moderate quality, 4–6 stars; or poor quality, 0–3 stars [32].

Statistical analysis

The Review Manager (RevMan) Version 5.3 software was used for data analysis. The pooled odds ratios (ORs) along with 95% confidence intervals (CIs) were used to calculate the risk of COVID-19 and associated outcomes in patients with PD and periodontium healthy patients. Heterogeneity was evaluated using the Chi-square test and the I^2 statistics. A Fixed-effects model was used for low/moderate heterogeneity

($I^2 \leq 50\%$), while a random effect model was applied for significant heterogeneity ($I^2 > 50\%$).

Sensitivity tests: due to the limited number of analyzed studies, no sensitivity tests were done.

Results

Study selection

Figure 1 presents the search strategy of the present study. The initial online searches yielded 3002 articles, of which 2050 duplicate records were excluded. The titles and abstracts of 952 articles were screened for eligibility, and 895 were found irrelevant and thus excluded. The full-texts of the remaining 57 articles were assessed for eligibility, and 35 were excluded for various reasons (Supplementary Table 2). Eventually, 22 studies were included in the present systematic review, and 12 were pooled into the meta-analysis.

General characteristics of the included studies

A total of 22 studies, involving 92,535 patients were included in the present systematic review; [22–25, 27–29, 33–47] of these, 12 studies were included in the meta-analysis [23, 25, 27, 28, 33, 35, 36, 40, 41, 43, 44, 46, 47]. Twelve of these studies were case–control studies, [22–25, 29, 34, 35, 37, 38, 42, 44, 46] three cohort studies, [27, 28, 39] and seven cross-sectional studies [33, 36, 40, 41, 43, 45, 47]. Six studies were conducted in India [23, 34, 38, 40, 42, 44] three in the UK [27, 28, 39], two in Brazil [35, 36] two in Saudi Arabia [33, 43], two in Qatar [22, 25], two in Turkey [46, 47] and one each in: the Netherlands, [29]. Spain, [45] Egypt, [41] Mexico, [37] and the USA [24]. The mean age of the participants ranged from 38.2 to 68.6 years, with almost equal representation of both genders. Diagnosis of COVID-19 was confirmed by PCR test in all of the included studies.

Periodontal parameters (Exposure)

Ascertainment of periodontal parameters were highly variable across the included studies. One or more of the following periodontal parameters were considered: number of missing teeth, pocket depth, gum bleeding, alveolar bone loss, and loose teeth (Table 1). Similarly, ascertainment of the exposure methods varied greatly across the included studies: clinical examination in nine studies, [23, 34–36, 38, 40, 42, 44, 46] self-reported in five studies, [27, 28, 37, 39, 41] and dental radiographs in eight studies [22, 24, 25, 29, 33, 43, 45, 47].

Outcome measures

Most of the studies reported on adverse outcomes of Covid-19. Ascertainment of the adverse outcomes




Fig. 1 Flowchart of the search strategy

of COVID-19 included one or more of the following: severity of symptoms, ICU admission, hospital admission, and mortality (Table 1). Nine studies [24, 26–28, 34, 37–39, 46] also reported on the risk of covid-19 in periodontitis patients.

Qualitative results

The majority of the studies (19 studies) found a significant association between PD and COVID-19 adverse outcomes (i.e., severity of symptoms, hospital admission, ICU and mortality). Conversely, three studies did not report a significant association between PD and COVID-19 adverse outcomes [27–29] except for mortality in one study [28]. Concerning susceptibility to COVID-19, nine studies [26–28, 34, 37–39, 46] reported on this outcome; six of these studies reported a significant association between having PD and the risk for COVID-19 infection, [24, 26, 34, 37, 38, 46] whereas the other three studies [27, 28, 39] didn’t confirm such results (Table 2).

Meta-analysis results

COVID-19 outcomes in PD versus healthy periodontium patients:

The pooled data showed a positive significant association between PD and the risk of adverse COVID-19 outcomes (Figs. 2, 3 and 4). Compared to patients with healthy periodontium, patients with PD showed a significantly higher risk of severe symptoms (OR=6.95, 95% CI: 2.24, 21.56, I²=92%, random-effect; P=0.0008)(Fig. 2), ICU admission (OR=3.15, 95% CI: 2.07, 4.79, I²=0.00%, fixed-effect; P=0.0001) (Fig. 3), and mortality (OR=1.92, 95% CI: 0.70, 5.32, I²=57%, random-effect; P=0.21) (Fig. 4).

COVID-19 outcomes by severity of PD (severe PD versus mild PD)

The results revealed a positive significant association between the severity of PD and severity of COVID-19 outcomes (Figs. 5, 6 and 7): severe symptoms (OR=3.25, 95% CI: 1.23, 8.59, I²=87%, random-effect; P=0.02) (Fig. 5); ICU admission (OR=3.38, 95% CI:

Table 1 General characteristics of the included studies

Author (country)	Study Design	No of Participants M/F (age)	Perio-related parameters assessment (Exposure)	COVID-19 Outcome parameters (Outcomes)	confounding factors
Anand et al., 2021 [22] (India)	Case-control	Total N= 150; Gender: 85M/65 F Cases: COVID-19 positive patients (n=79) (Mean age: 43.34) Controls: COVID-19 negative (n=71) (Mean age: 38.24)	Pl, CI, PD, GB, tooth mobility, recession, CAL (clinical exam)	COVID-19 infection risk	Tobacco use, age, systemic diseases, oral hygiene practices Diabetes Hypertension Neoplasma
Marouf et al., 2021 [25] (Qatar)	Case-control	Total N= 568; Gender: 310/258 Cases: COVID-19 patients with complications (n=40) (mean age: 53.5 ys) Control: COVID-19 patients without complications (n=528) (mean age: 41.5)	Interdental bone loss Periodontal condition (radiographs)	Risk of COVID-19 infection, COVID-19 complications: ICU, death, assisted ventilation	age, sex, smoking, BMI, diabetes and co-morbidities.
Larvin et al., 2020 [29] (UK biobank)	Retrospective cohort	Total N: 13,253 biobank (1,616 positive covid and 11,637 negative) Gender: 6451/6802 Age: 68.55 years Cases: patients with PD (n= 2100) Controls: patients with no PD (n= 11,153)	Painful/bleeding gums Loose teeth (Self-reported)	Risk of COVID-19, COVID-19 complications -Hospital admission - mortality	age, sex, ethnicity, income, smoking, BMI, cancer, diabetes and CVD
Gupta et al., 2021 [23] (India)	Case-control	Total n= 82; gender: 48/34 Cases: 55 COVID-19 patients with PD (n= 55) Controls: 27 COVID-19 patients with no PD (n=27) Mean Ages: 34.44 to 62.94	Pocket depth, CAL, BOP Gingivitis, Periodontitis(stage I, stage II, stage III, stage IV) and Recession (Clinical exam)	pneumonia, death/survival, - hospital admission-assisted ventilation	age, sex, smoking habits , diabetes, hypertension, pulmonary disease, chronic kidney disease, cancer, CAD, obesity
Holt et al., 2021 [40] (UK)	Cohort study	15227 participants (446 COVID-19 cases) Gender: 4597/10630 Age: 59.4 years	Painful/bleeding gums Loose teeth (Self-reported)	Risk of COVID-19 infection	age, sex, alcohol and smoking habits, ethnicity, co-morbidities BMI
Donders et al., 2022 [31] (Netherlands)	Case-control study	Total : N=133 Gender: 61/72 Age: 61.7 years G1: Mild COVID-19 (n=82) G2: moderate (n=30) G3: Severe COVID-19 (n=21)	-Periodontitis -tooth loss (Radiographs)	Severity of COVID = -Mild, -moderate and, Severe (ICU/ death, ICU)	age, sex, smoking, diabetes, CVD, kidney diseases

Table 1 (continued)

Author (country)	Study Design	No of Participants M/F (age)	Perio-related parameters assessment (Exposure)	COVID-19 Outcome parameters (Outcomes)	confounding factors
Larvin et al., 2021 [30] (UK)	Retrospective cohort study periodontal disease in obesity on COVID-19	Total n= 58,897; 14,466 (24.6%) COVID-19 positive PD patients: n= 9332 No PD: n= 49,565 age: 56.83 y 53% females	bleeding gums, painful gums, and loose teeth (Self-reported)	risk of COVID-19 infection, hospital admission, and mortality	age, sex, ethnicity, income, smoking, BMI, cancer, diabetes and CVD
Costa et al., 2022 [37] (Brazil)	Prospective Cross-sectional study	Total N=128 COVID-19 positive Gender 68/60 Age: 58.7y	DMFT index, Probing depth (PD), clinical attachment level (CAL), bleeding on probing (Clinically)	-Severity (mild, moderate, severe, or critical) -Hospital admission -discharge or death -ventilation	Patient age Comorbidities(diabetes, hypertension, and obesity)
Kaur et al., 2022 [42] (India)	Case-control	N=116 subjects suffering from COVID-19 Gender 70/46 Two groups G1: Moderate COVID-19 cases (G2: Mild COVID-19	Periodontitis(Stage 0-1) Periodontitis(Stage 2-3) (Clinically)	Severity of COVID = -Mild, -moderate	Smoking, diabetes, age, comorbidities, BMI
Mishra et al., 2022 [44] (India)	Case-control	N= 294 subject with Covid Gender 154/140 COVID-19 cases were divided into: G1 : Mild cases (163) Moderate cases (83) Severe cases (48)	Periodontitis (Stage 1-11) Peri-odontitis (Stage III-IV) PI, CI, PD, GB, tooth mobility, recession, CAL (Clinically)	Severity of COVID = -Mild, -moderate, severe	Smoking, diabetes, age, gender, BMI, oral hygiene practices
Kamel et al., 2021 [24] (Egypt)	Cross-sectional	N= 308 Age: 18-55 years No gender reported	Gum bleeding, tooth loss, gum pain (Self-reported)	Severity of COVID-19 (severe and mild) (dyspnea, low O2 saturation, hospitalization),	Participants with known risk factors and comorbidities that could affect COVID-19 severity were excluded.
Sirin & Ozcelik, 2021 [47] (Turkey)	Comparative Cross-sectional	N= 137 Age: 20-65 yrs Gender: 71/66	Alveolar bone loss, tooth loss, Dental implants Apical periodontitis (Radiographs)	Severity of COVID-19	NA
Koppolu et al., 2023 [43] (KSA)	Retrospective comparative Cross-sectional	N= 196 Age: 18-60 yrs Gender: 62/36	Alveolar bone loss (Radiographs)	Severity of COVID-19 (COVID negative, Home isolation, Hospital without ventilator and ICU, Ventilation, ICU, Death)	Age, sex, smoking, comorbidity.
Sari et al., 2023 [46] (Turkey)	Case-control	N= 80 (50% had COVID) Age: 23-64 years Gender: 38/42	PD, CAL, PI, GI, BOP (Clinically)	Prevalence of COVID-19 infection among patients with periodontitis.	Age, gender, smoking, BMI.

Table 1 (continued)

Author (country)	Study Design	No of Participants M/F (age)	Perio-related parameters assessment (Exposure)	COVID-19 Outcome parameters (Outcomes)	confounding factors
Said et al., 2023 [27] (Qatar)	Case-control	N= 1325 (71 suffered severe COVID-19 complications resulting in ICU admission, mechanical ventilation and/or death). Age: > 18 years Gender: 577/748	Inter-dental bone loss in posterior teeth (Radiographs)	COVID-19 complications (any complication, ICU admission, ventilation and death)	Age, sex, diabetes, smoking habits, BMI and comorbidities
Alnomay et al., 2023 [35] (KSA)	Retrospective cross-sectional	N= 188 Age: ≥ 18 years Gender: 81/107	Inter-dental bone loss in posterior teeth (Radiographs)	Severity of COVID-19	Age, gender, smoking, diabetes, hypertension, obesity, and comorbidities
Guardado-Luevanos et al., 2022 [38] (Mexico)	Case-control	N= 234 Age: 20-75 years Gender: 96/138	periodontal disease (Self-reported)	Prevalence of COVID-19 infection among patients with self-reported periodontitis.	Age, sex, smoking, and comorbidities.
Gujar et al., 2022 [39] (India)	Case-control	N= 150 (79 were categorized as cases) Age: ≥ 18 years No gender reported	Plaque scores, calculus scores, tooth mobility, gingival bleeding, probing depth (PD), recession (REC), and clinical attachment level (CAL) (Clinically)	Prevalence of COVID-19 infection among patients with periodontitis.	Age, sex, smoking, oral hygiene habits, and comorbidities.
Wadhwa et al., 2022 [26] (USA)	Case-control	N= 387 COVID-19 cases, 387 control Age: 47.1 years Gender: 260/514	Missing teeth Alveolar bone (Radiographs)	-Risk of Covid-19 -Severity of Covid-19	Age, gender, smoking, race, ethnicity
Poyato-Borrego et al., 2023 [45] (Spain)	Retrospective comparative Cross-sectional	N= 52 Age: 52.3 Gender: 30/22	Alveolar bone loss (Radiographs)	Severity of COVID-19; mild/moderate disease (MMG) and severe/critical disease (SCG)	NA
Kalsi et al., 2023 [41] (India)	Cross-sectional	N= 44 Age: 20-50 years Gender: 23/21	Probing pocket depth (PPD), clinical attachment level (CAL), bleeding on probing (BOP), periodontal inflamed surface area (PISA) (Clinically)	Severity of COVID-19	NA
Bemquerer et al., 2023 [36] (Brazil)	Case-control	N= 99 cases; 182 control Age: 16-74 yrs cases; 20-74 yrs control Gender: 52/47 cases; 96/86 control	Plaque index, probing depth (PD), clinical attachment level (CAL), and bleeding on probing (BOP). (Clinically)	Severity of COVID-19 (use of oxygen, hospitalization, admission to the intensive care unit (ICU), admission to the semi-intensive care unit (SICU), use of mechanical ventilation, length of hospitalization, length of ICU admission, length of mechanical ventilation, and death)	Age, sex, schooling, family income, toothbrushing, flossing, smoking, and BMI.

M male, F female, PD periodontal disease, CAD coronary artery disease, CVD Cardiovascular disease, MM not available

Table 2 Summary of the main outcomes

Study	Main outcome
Anand et al., 2021 [22]	There was a highly significant association between some PD parameters and the risk of COVID-19 : plaque index (OR 7.01; 95% CI, 1.83 -26.94), gingival inflammation (OR, 17.65; 95% CI, 5.95 - 52.37), CAL (OR, 8.46; 95% CI, 3.47 to 20.63), periodontitis (OR, 11.75; 95% CI, 3.89 to 35.49).yet, there were no significant differences between the two groups in terms of missing teeth, carious teeth, and calculus scores.
Marouf et al., 2021 [25]	A significant association between periodontitis and COVID-19 complications including death (OR = 8.81, 95% CI 1.00–77.7), ICU admission (OR = 3.54, 95% CI 1.39–9.05) and the need for assisted ventilation (OR = 4.57, 95% CI 1.19–17.4). Furthermore, white blood cells, D-dimer and CRP levels were significantly higher in COVID-19 patients with PD.
Larvin et al 2020 [29]	There was no any association between painful/bleeding gum and the risk of COVID-19 (OR = 1.10, 95% CI = 0.72–1.69) or risk of hospital admission (OR = 0.90, 95% CI = 0.59–1.37). However, there was a significant association between PD and the mortality rate; participants with painful/bleeding gum had twofold higher mortality rate than those with healthy gum (OR = 1.71, 95% CI = 1.05–2.72). There was no a significant association between loose teeth with COVID-19 parameters: risk of COVID-19 (OR = 1.15, 95% CI = 0.84–1.59); hospital admission (OR = 1.55, 95% CI = 0.87–2.77); or mortality (OR = 1.85; 95% CI = 0.92–2.72)
Gupta et al. 2021 [23]	There is a significant association between PD and COVID-19 - outcomes. Higher severity of PD was associated with 7.45 odds of requiring assisted ventilation, 36.52 odds of hospital admission, 14.58 odds of death, and 4.42 odds of COVID-19 -related pneumonia
Holt et al. 2021 [40]	The results showed a significant increase in the risk of COVID-19 in PD subjects (OR: 1.20). However, after controlling all potential confounding factors, no significant association was observed between the two conditions.
Donders et al 2022 [31]	The results showed a significant association between PD (alveolar bone loss [OR: 5.60; 95%CI: 1.21; 25.99; $P = 0.028$] and tooth loss [OR: 1.04; 95%CI: 1.00; 1.09; $P = 0.047$]) with severity of COVID-19. However, such associations disappeared after adjusting for all potential confounders (P more than 0.05).
Larvin et al 2021 [30]	The risk for COVID-19 infection was not associated with periodontal disease across the 3 BMI categories: normal weight (OR, 0.97; 95% CI, 0.88 to 1.07), overweight (OR, 1.06; 95% CI, 0.98 to 1.15), and obese (OR, 1.08; 95% CI, 0.99 to 1.17). The risk of hospital admission for people with periodontal disease was 38% higher in participants who were overweight (HR, 1.38;95% CI, 1.02 to 1.87) and 124% higher in those who were obese (HR, 2.24; 95% CI, 1.66 to 3.03) compared to those of normal weigh. In addition, for participants with obesity, the mortality rate was much higher (hazard ratio, 3.11; 95% CI, 1.91 to 5.06) in participants with periodontal disease than those with healthy periodontium.
Costa et al., 2022 [37]	Periodontitis was significantly associated with ICU admission [IRR = 1.44 (95%CI = 1.07–1.95); $p = 0.017$], critical symptoms [IRR = 2.56 (95%CI = 1.44–4.55); $p = 0.001$], and risk of death [IRR = 2.05 (95%CI = 1.12–3.76); $p = 0.020$]
Kaur et al 2022 [42]	The risk of severe periodontal disease was 6.32 times higher in COVID-19 patients with severe symptoms ($P = 0.024$) as compared to mild COVID cases.
Mishra et al 2022 [44]	There was a significant association between periodontitis and severity of COVID-19 symptoms. The adjusted odds ratio (OR) of having severe COVID-19 in periodontitis patients was 2.8133 (0.4077–19.7523 at 95% CI, $p = 0.004$).
Kamel et al. 2021 [24]	There was a significant association between poor oral health and the severity of COVID-19. Additionally, poor oral health was significantly associated with delayed recovery time and higher CRP values.
Sirin & Ozcelik 2021 [47]	There was a significant association between dental disease including alveolar bone loss and severity of COVID-19 complications. Individuals with severe dental diseases (including alveolar bone loss) had more severe COVID-19 symptoms and higher mortality rate than those with no/mild dental diseases.
Koppolu et al., 2023 [43]	There was a statistically significant link between gingivitis and periodontitis. The majority of mild gingivitis cases (63%) was associated with the COVID-19-negative group, whereas the majority of severe gingivitis groups (85.7%) was associated with the COVID-19-positive group ($\chi^2 = 9.94$; $P = 0.007$). Similarly, the majority of Stage 1 periodontitis (62.9%) was associated with COVID-19-negative participants, whereas the majority of Stage 4 periodontitis ($P = 0.007$) was associated with COVID-19-positive groups ($\chi^2 = 22.51$; $P = 0.047$).
Sari et al., 2023 [46]	A higher prevalence of periodontitis compared with gingivitis and periodontal health was detected in the test group ($p = 0.003$). In line with this, all clinical periodontal parameters related to periodontal disease severity were higher in the test group than in the control group. In particular, GI, BOP (%), PD, CAL, and the number of missing teeth were statistically significantly higher in the test group than in the control group ($p < 0.05$), except PI, which did not reach statistical significance ($p = 0.052$).
Said et al., 2023 [27]	Risk analysis of COVID-19 complications revealed that while periodontitis stage 2–4 (regardless of treatment) was associated with higher risk of complications (i.e. need for mechanical ventilation [AOR = 3.32, 95% CI 1.10–10.08, $p = 0.034$]), subjects with treated periodontitis had a lower risk than the non-treated ones. Adjusted OR analysis comparing treated and non-treated periodontitis (stages 2–4) revealed that treated patients were at lower risk of complications; however, this was not statistically significant.
Alnomay et al., 2023 [35]	Patients with periodontitis were 3-times 182 more likely to have COVID-19 complications than those without periodontitis ($p = 0.025$).
Guardado-Luevanos et al., 2022 [38]	A statistically significant difference (p value < 0.001) was observed, showing that positive self-RPD ($n = 95$, 85.1%) was often higher in SARS-CoV-2-positive individuals than in the controls ($n = 66$, 56.4%), with an OR of 3.3 (1.8–6.0) for SARS-CoV-2 infection in people with self-reported periodontal disease.

Table 2 (continued)

Study	Main outcome
Gujar et al., 2022 [39]	Participants with COVID-19 had significantly higher mean values of plaque scores, number of mobile teeth, gingival bleeding scores, PD, REC, and CAL compared to the controls. The mean percentages of inter-proximal sites with PD ≥ 4 mm, PD ≥ 5 mm, CAL ≥ 3 mm, CAL ≥ 4 mm, and CAL ≥ 6 mm were also significantly higher in the case group than in the control group.
Wadhwa et al. 2022 [26]	Covid-19 infected patients had significantly greater alveolar bone loss and missing teeth than controls. Additionally, missing teeth and bone loss were associated with more hospitalization.
Poyato-Borrego et al., 2023 [45]	A significant association has been found between the severity of COVID-19 and the periodontal disease ($p = 0.04$).
Kalsi et al., 2023 [41]	The percentage of patients suffering from severe COVID disease was least (8.3%) in stage 1 of periodontitis and was highest (62.5%) in stage 4 of periodontitis. Whereas out of eight patients tested positive in stage 1 of periodontitis, 7 had only mild disease. Out of total of 8 patients with stage 4 category of periodontitis, two remain uninfected, five developed severe form of COVID, and only one patient had moderate COVID disease.
Bemquerer et al., 2023 [36]	In the COVID-19 group, plaque index was worse in patients with periodontitis than in patients without periodontitis ($p < 0.001$). Among individuals with COVID-19, periodontitis was associated with more hospitalization ($p = 0.009$), more days in the ICU ($p = 0.042$), admission to the SICU ($p = 0.047$), and higher need for oxygen therapy ($p = 0.042$). However, there was no difference in total hospitalization length, need for ICU, need for mechanical ventilation, or death between individuals with and without periodontitis in the COVID-19 group. Individuals with periodontitis were 1.13 times more likely to be hospitalized than individuals without periodontitis ($CI = 1.01-1.26$; $p = 0.028$).

PD periodontal disease, OR odds ratio, CI confidence interval, MR Mendelian randomization, IVW Inverse-variance weighted, CRP C-reactive protein




Fig. 2 Meta-analysis of the association between periodontal disease (PD) and severe COVID-19 symptoms




Fig. 3 Meta-analysis of the association between periodontal disease (PD) and ICU admissions




Fig. 4 Meta-analysis of the association between periodontal disease (PD) and mortality




Fig. 5 Meta-analysis of the association between severe periodontal disease (PD) and COVID-19 symptoms




Fig. 6 Meta-analysis of the association between severe periodontal disease (PD) and ICU admission




Fig. 7 Meta-analysis of the association between severe periodontal disease (PD) and mortality rate




Fig. 8 Meta-analysis of COVID-19 risk in relation to periodontal health status (PD vs. healthy)

1.82, 6.26, I² = 0.00%, fixed-effect; P = 0.0001) (Fig. 6), and mortality rate (OR = 5.35, 95% CI: 2.75, 10.42, I² = 48%, fixed-effect; P = 0.00001) (Fig. 7).

Risk of COVID-19 in PD patients versus healthy periodontium patients:

The pooled three studies revealed a higher risk of COVID-19 infection in periodontitis patients (OR = 1.58, 95% CI: 0.89, 2.79, I² = 57%, P = 0.12), although the result was not statistically significant (Fig. 8).

Publication bias

The funnel plots for publication bias of the include studies for all categories are presented in the Supplementary file 2.

Generally, there was not a noticeable publication bias among the studies, except Fig. 1 in the Supplementary file 2 which showed slight publication bias for the association between periodontal disease (PD) and severe COVID-19 symptoms.

Quality assessment

The results of the quality assessment- based on NOS- are presented in Table 3. Overall, the included studies revealed relatively good quality ranging from five to nine stars. Fourteen studies [22–25, 27–29, 34, 36–38, 40, 43–45] were of high quality, while seven studies were of moderate quality [33, 35, 39, 41, 42, 46, 47]. The most frequent methodological shortcomings were related to the self-report ascertainment of the exposure and bias in selection of cases/controls (Table 3).

Table 3 NOS-based quality analysis of the included studies

Study	Selection	Comparability	Exposure/Outcome	Overall
Anand et al., 2021 [22]	****	**	***	Low
Marouf et al., 2021 [25]	****	**	***	Low
Larvin et al 2020 [29]	***	**	***	Low
Gupta et al. 2021 [23]	***	**	**	Low
Holt et al. 2021 [40]	**	*	***	Moderate
Donders et al 2022 [31]	***	**	***	Low
Larvin et al 2021 [30]	***	**	**	Low
Costa 2022 [37]	***	**	**	Low
Kaur et al 2022 [42]	**	*	***	Moderate
Mishra et al 2022 [44]	***	**	**	Low
Kamel et al. 2021 [24]	***	*	*	Moderate
Sirin & Ozcelik 2021 [47]	**	*	**	Moderate
Koppolu et al., 2023 [43]	**	**	***	Low
Sari et al., 2023 [46]	**	*	**	Moderate
Said et al., 2023 [27]	****	**	***	Low
Alnomay et al., 2023 [35]	**	**	**	Moderate
Guardado-Luevanos et al., 2022 [38]	**	**	***	Low
Gujar et al., 2022 [39]	**	**	***	Low
Wadhwa et al. 2022 [26]	***	**	***	Low
Poyato-Borrego et al., 2023 [45]	****	**	***	Low
Kalsi et al., 2023 [41]	***	**	***	Low
Bemquerer et al., 2023 [36]	**	**	**	Moderate

Discussion

Medical literature has linked PD to the risk and severity of COVID-19. Hence, the present systematic review and meta-analysis sought to answer the following focused question: Does PD influence the risk and severity of COVID-19? Qualitatively, most of the included studies reported significant association between PD and COVID-19 severity. However, three studies [27–29] failed to replicate these results. Quantitatively the pooled data found a significant positive association between PD and the risk and adverse outcomes of COVID-19 such as severe symptoms, ICU admission. Additionally, severe PD was significantly associated with higher risk of severe COVID-19 symptoms ($P=0.02$), ICU admission ($P=0.0001$), and mortality rate ($P=0.00001$) compared to mild PD. Indeed, the results revealed that patients with PD have significantly 54% higher risk to getting COVID-19 infection. However, these findings should be interpreted with caution owing to the heterogeneity among the included studies as well as some methodological limitations, discussed in the following sections.

The findings of the present systematic review support the results of previous systematic reviews [30, 31]. Nevertheless, although the results of the current study are interesting, the mechanism(s) by which periodontitis aggravate(s) COVID-19 adverse outcomes still unclear so far. However, many theories have been suggested, and deserve discussion. With regard to one of the key findings of the current study: association of PD and COVID-19 adverse outcomes, one possible explanation is related to the expression of angiotensin converting enzyme 2 (the well-known receptor for SARS-CoV-2) by the inspired periopathogenes. This subsequently leads to production of inflammatory cytokines such as IL-6 and IL-8 in the lower respiratory tract, thus aggravating the response [48]. Further, periopathogenes have been reported to enhance the virulence of SARS-CoV-2 by cleaving its S glycoproteins, a matter that exacerbates COVID-19 complications [49]. Of utmost important, the chronic inflammatory nature of periodontitis may play a role through triggering systemic inflammation, which aggravates the inflammatory response in context of many disease processes, and COVID-19 wouldn't be an exception. A recent study reported existence of periopathogenes in the metagenome of patients severely infected with SARS-CoV-2: Mainly high reads for *Prevotella* (493 reads), *Staphylococcus* (1,659 reads) and *Fusobacterium* (463 reads) were discovered [50]. Indeed, the potential role of PD in pulmonary infections (and diseases) has long been investigated and well documented in the literature [16, 51–53]. Mounting evidence from systematic reviews and meta-analyses found a significant association

between PD and exacerbation of respiratory conditions, mainly pneumonia and COPD [14, 17].

The significant association between PD and the risk of COVID-19 is another key and pivotal finding of the current study: patients with PD were 54% at higher risk of COVID-19 acquisition than people with healthy periodontium. Essentially, the oral cavity, including gingival pocket epithelium, has been reported to be potentially high risk for SARS-CoV-2 infectious susceptibility, mediated by expression of angiotensin converting enzyme 2 [54, 55]. These results might be explained by the role of periopathogenic bacteria in initiating the expression of angiotensin converting enzyme, as already discussed above. A very recent evidence suggests that periodontal pockets and decayed teeth serve as reservoirs for SARS-CoV-2, making people more prone to COVID-19 [56]. In addition to being hiding areas, where the antibacterial action of the saliva and mouth rinses is not effective, the periodontally involved pockets have higher surface area than that of normal gingival sulcus, providing more opportunities for SARS-CoV-2 to bind and eventually infect more enzyme-expressing cells, in addition to acting as reservoirs for continuing infection, or recurring, or complicating the current infection leading to a more severe disease. However, these are still hypotheses and more research are required to elucidate these phenomena.

Worthy to note that the positive association of PD with the outcomes of COVID-19 was supported by all of the included studies except two studies by Larvin et al., which failed to do so [27, 28]. It must be acknowledged that these two studies involved relatively large sample sizes of COVID-19 patients. However, the exposure (periodontal status) was self-reported, a matter which raises a lot of doubts and reduces reliability of these two studies. Obviously, it is a clear shortcoming, and therefore the results might have not accurately reflected the periodontal status of the participants, and thus might have biased the results. Apart from the fact that the number of present and/or the missing teeth can be self-reported to a large extent of reliability, the other periodontal health outcomes and parameters cannot be reported by the patients precisely [57]. This, in turn, may explain the different results obtained by these two studies, a matter that must be acknowledged too. By contrast, except for Kamel et al. study, all other studies, which reported positive association of PD with the outcomes of COVID-19 ascertained the periodontal parameters objectively, either clinically or using radiographs.

Undoubtedly, the quality of the individual studies is a very determining factor that influences the quality of the overall evidence of any meta-analysis [58]. For this purpose, two reviewers evaluated independently the

quality of all included studies using NOS, a very valid risk assessment tool. The results revealed relatively good quality of the included studies. Selection bias and self-reporting of the exposure (periodontal parameters) were the most frequently shortcomings, which cause biased results: recapitulating and reminding Larvin et al. studies mentioned above.

The present systematic review has many points that add to its strength and should be recognized. First, the study included a good number of studies with a relatively large sample size. Second, the studies were conducted in different geographical areas representing the world and thus substantiating its external validity. However, there are few limitations that should be highlighted. The main limitation is the heterogeneity among the included studies in many respects including: study design, the assessed periodontal parameters (exposure), COVID-19 parameters (outcome), setting of the included patients (hospitalized vs non-hospitalized patients), age of the participants, and many other confounders. The differences in methods of measurements of periodontal parameters represent a marked heterogeneity, being self-reported in a few studies, using radiographs in some studies, and clinical examination in the other studies. In particular, the self-reporting of periodontal status is an evident drawback of the present study that might have caused bias in the results. Furthermore, missing of some of the numerical data was an obvious obstacle that hindered including all studies in the quantitative analysis.

Conclusion

In conclusion, the present systematic review and meta-analysis suggests a significant association between poor periodontal health and poor COVID-19 outcomes. However, the results should be interpreted with caution given the marked heterogeneity across the included studies along with some methodological limitations in some of these studies. Hence, further large-scale prospective cohort studies with standardized methodologies are highly required to further elucidate the potential association between periodontal diseases and the risk of poor COVID-19 outcomes.

Abbreviations

COVID-19	Coronavirus Disease 2019
PD	Periodontitis
ICU	Intensive Care Unit
CI	Confidence Interval
NOS	Newcastle Ottawa Scale
OR	Odds Ratio

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s12903-023-03378-0>.

Additional file 1: Table S1. Databases. Applied search strategy, and numbers of retrieved studies. **Table S2.** List of excluded studies and reason for exclusion.

Additional file 2: Figure S1. Funnel plot for the included studies of the association between periodontal disease (PD) and severe COVID-19 symptoms. **Figure 2.** Funnel plot for the included studies of the association between periodontal disease (PD) and ICU admissions. **Figure S3.** Funnel plot for the included studies of the association between periodontal disease (PD) and mortality. **Figure S4.** Funnel plot for the included studies of the association between severe periodontal disease (PD) and COVID-19 symptoms. **Figure S5.** Funnel plot for the included studies of the association between severe periodontal disease (PD) and ICU admission. **Figure S6.** Funnel plot for the included studies of the association between severe periodontal disease (PD) and mortality rate. **Figure S7.** Funnel plot for the included studies of COVID-19 risk in relation to periodontal health status (PD vs. healthy).

Acknowledgements

Open Access funding was provided by Qatar National Library.

Authors' contributions

Sadeq Al-Maweri: study concept, search strategy, drafting the manuscript; Mohammed Nasser Alhaji: data extraction, quality appraisal, data analysis, drafting the manuscript; Esam Halboub: concept of the study, critically revised and edited the paper; Faleh Tamimi: data extraction, quality appraisal, drafting the manuscript; Nosizana Mohd Salleh: concept of the study, critically revised and edited the paper; Mohammed Sultan Al-Ak'hali: concept of the study, critically revised and edited the paper; Saba Kassim, Saleem Abdulrab, Lamyia Anweigi, and Marwan Mansoor Ali Mohammed, data curation, critically revised and edited the paper. All authors approved the final version.

Funding

No fund was received for this study.

Availability of data and materials

All data generated or analyzed during this study are included in this published article.

Declarations

Ethics approval and consent to participate

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Consent for publication

Not applicable.

Author details

¹College of Dental Medicine, QU Health, Qatar University, Doha, Qatar. ²Department of Restorative Dentistry, Faculty of Dentistry, University of Malaya, Federal Territory of Kuala Lumpur, Kuala Lumpur, Malaysia. ³Department of Prosthodontics, Faculty of Dentistry, Tamar University, Dhamar, Yemen. ⁴Department of Maxillofacial Surgery and Diagnostic Sciences, College of Dentistry, Jazan University, Jazan, Saudi Arabia. ⁵Department of Preventive Dental Sciences, College of Dentistry, Jazan University, Jazan, Saudi Arabia. ⁶Department of Preventive Dental Sciences, College of Dentistry, Taibah University, Al-Madinah Al-Munawwarah, Saudi Arabia. ⁷Al Khor Health Center, Primary Health Care Corporation, Doha, Qatar. ⁸Department of Oral and Craniofacial Health Sciences, College of Dental Medicine, University of Sharjah, Sharjah, United Arab Emirates.

Received: 21 June 2023 Accepted: 31 August 2023
Published online: 09 September 2023

References

- DeMartino JK, Swallow E, Goldschmidt D, Yang K, Viola M, Radtke T, Kirson N. Direct health care costs associated with COVID-19 in the United States. *J Manag Care Spec Pharm*. 2022;28(9):936–47.
- Prezant DJ, Lancet EA, Zeig-Owens R, Lai PH, Appel D, Webber MP, Braun J, Hall CB, Asaada G, Kaufman B, et al. System impacts of the COVID-19 pandemic on New York City's emergency medical services. *J Am Coll Emerg Physicians Open*. 2020;1(6):1205–13.
- Al-Maweri SA, Halboub E, Warnakulasuriya S. Impact of COVID-19 on the early detection of oral cancer: A special emphasis on high risk populations. *Oral Oncol*. 2020;106: 104760.
- Zhou F, Yu T, Du R, Fan G, Fan Y, Liu X, Xiang J, Wang Y, Song B, Gu X, et al. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet*. 2020;395(10229):1054–62.
- Richardson S, Hirsch JS, Narasimhan M, Crawford JM, McGinn T, Davidson KW, Barnaby DP, Becker LB, Chelico JD, Cohen SL, et al. Presenting Characteristics, Comorbidities, and Outcomes Among 5700 Patients Hospitalized With COVID-19 in the New York City Area. *JAMA*. 2020;323(20):2052–9.
- Pujhari S, Paul S, Ahluwalia J, Rasgon JL. Clotting disorder in severe acute respiratory syndrome coronavirus 2. *Rev Med Virol*. 2021;31(3): e2177.
- Sinha P, Matthay MA, Calfee CS. Is a "Cytokine Storm" Relevant to COVID-19? *JAMA Intern Med*. 2020;180(9):1152–4.
- Liu J, Virani SS, Alam M, Denktas AE, Hamzeh I, Khalid U. Coronavirus disease-19 and cardiovascular disease: A risk factor or a risk marker? *Rev Med Virol*. 2021;31(3): e2172.
- Papapanou PN, Sanz M, Buduneli N, Dietrich T, Feres M, Fine DH, Flemmig TF, Garcia R, Giannobile WV, Graziani F, et al. Periodontitis: Consensus report of workgroup 2 of the 2017 World Workshop on the Classification of Periodontal and Peri-Implant Diseases and Conditions. *J Clin Periodontol*. 2018;45(Suppl 20):S162–s170.
- Petersen PE, Ogawa H. The global burden of periodontal disease: towards integration with chronic disease prevention and control. *Periodontol* 2000. 2012, 60(1):15–39.
- Al-Maweri SA, Ibraheem WI, Al-Ak'hali MS, Shamala A, Halboub E, Alhajj MN. Association of periodontitis and tooth loss with liver cancer: A systematic review. *Crit Rev Oncol Hematol*. 2021;159: 103221.
- Alakhali MS, Al-Maweri SA, Al-Shamiri HM, Al-Haddad K, Halboub E. The potential association between periodontitis and non-alcoholic fatty liver disease: a systematic review. *Clin Oral Investig*. 2018;22(9):2965–74.
- Dhaifullah E, Al-Maweri SA, Koppolu P, Elkhat E, Mostafa D, Mahgoub M. Body mass index and periodontal health status among young Saudi adults: a cross-sectional study. *Ann Saudi Med*. 2019;39(6):433–40.
- Kelly N, Winning L, Irwin C, Lundy FT, Linden D, McGarvey L, Linden GJ, El Karim IA. Periodontal status and chronic obstructive pulmonary disease (COPD) exacerbations: a systematic review. *BMC Oral Health*. 2021;21(1):425.
- Sanz M, Marco Del Castillo A, Jepsen S, Gonzalez-Juanatey JR, D'Aiuto F, Bouchard P, Chapple I, Dietrich T, Gotsman I, Graziani F, et al. Periodontitis and cardiovascular diseases: Consensus report. *J Clin Periodontol*. 2020;47(3):268–88.
- Scannapieco FA, Bush RB, Paju S. Associations between periodontal disease and risk for nosocomial bacterial pneumonia and chronic obstructive pulmonary disease. A systematic review *Ann Periodontol*. 2003;8(1):54–69.
- Wu Z, Xiao C, Chen F, Wang Y, Guo Z. Pulmonary disease and periodontal health: a meta-analysis. *Sleep Breath*. 2022.
- Hajishengallis G. Periodontitis: from microbial immune subversion to systemic inflammation. *Nat Rev Immunol*. 2015;15(1):30–44.
- Tamimi F, Altigani S, Sanz M. Periodontitis and coronavirus disease 2019. *Periodontol* 2000. 2022, 89(1):207–214.
- Sahni V, Gupta S. COVID-19 & Periodontitis: The cytokine connection. *Med Hypotheses*. 2020;144: 109908.
- Herrera D, Serrano J, Roldán S, Sanz M. Is the oral cavity relevant in SARS-CoV-2 pandemic? *Clin Oral Invest*. 2020;24(8):2925–30.
- Marouf N, Cai W, Said KN, Daas H, Diab H, Chinta VR, Hssain AA, Nicolau B, Sanz M, Tamimi F. Association between periodontitis and severity of COVID-19 infection: a case-control study. *J Clin Periodontol*. 2021;48(4):483–91.
- Gupta S, Mohindra R, Singla M, Khara S, Sahni V, Kanta P, Soni RK, Kumar A, Gauba K, Goyal K, et al. The clinical association between Periodontitis and COVID-19. *Clin Oral Invest*. 2022;26(2):1361–74.
- Wadhwa S, Dave S, Daily ML, Nardone A, Li R, Rosario J, Cantos A, Shah J, Lu HH, McMahon DJ, et al. The role of oral health in the acquisition and severity of SARS-CoV-2: a retrospective chart review. *Saudi Dent J*. 2022;34(7):596–603.
- Said KN, Al-Momani AM, Almaseeh JA, Marouf N, Shatta A, Al-Abdulla J, Alaji S, Daas H, Tharupeedikayil SS, Chinta VR, et al. Association of periodontal therapy, with inflammatory biomarkers and complications in COVID-19 patients: a case control study. *Clin Oral Investig*. 2022;26(11):6721–32.
- Wang Y, Deng H, Pan Y, Jin L, Hu R, Lu Y, Deng W, Sun W, Chen C, Shen X, et al. Periodontal disease increases the host susceptibility to COVID-19 and its severity: a Mendelian randomization study. *J Transl Med*. 2021;19(1):528.
- Larvin H, Wilmott S, Kang J, Aggarwal VR, Pavitt S, Wu J. Additive Effect of Periodontal Disease and Obesity on COVID-19 Outcomes. *J Dent Res*. 2021;100(11):1228–35.
- Larvin H, Wilmott S, Wu J, Kang J. The Impact of Periodontal Disease on Hospital Admission and Mortality During COVID-19 Pandemic. *Front Med (Lausanne)*. 2020;7: 604980.
- Donders H, van der Sleen J, Kleinbergen Y, Su N, de Lange J, Loos B. Alveolar bone loss and tooth loss are associated with COVID-19 severity but are not independent risk factors. An explorative study. *J Oral Maxillofac Surg*. 2022, 5:100223.
- Espinoza-Espinoza DAK, Dulanto-Vargas JA, Cáceres-LaTorre OA, Lamas-Castillo FE, Flores-Mir C, Cervantes-Ganoza LA, López-Gurreonero C, Ladera-Castañeda MI, Cayo-Rojas CF. Association between periodontal disease and the risk of COVID-19 complications and mortality: a systematic review. *J Int Soc Prev Community Dent*. 2021;11(6):626–38.
- Baima G, Marruganti C, Sanz M, Aimetti M, Romandini M. Periodontitis and COVID-19: biological mechanisms and meta-analyses of epidemiological evidence. *J Dent Res*. 2022;101(12):1430–40.
- Stang A. Critical evaluation of the Newcastle-Ottawa scale for the assessment of the quality of nonrandomized studies in meta-analyses. *Eur J Epidemiol*. 2010;25(9):603–5.
- Alnomay N, Alolayan L, Aljohani R, Almashouf R, Alharbi G. Association between periodontitis and COVID-19 severity in a tertiary hospital: A retrospective cohort study. *Saudi Dent J*. 2022;34(7):623–8.
- Anand PS, Jadhav P, Kamath KP, Kumar SR, Vijayalaxmi S, Anil S. A case-control study on the association between periodontitis and coronavirus disease (COVID-19). *J Periodontol*. 2022;93(4):584–90.
- Bemquerer LM, Oliveira SR, de Arruda JAA, Costa FPD, Miguita L, Bemquerer ALM, et al. Clinical, immunological, and microbiological analysis of the association between periodontitis and COVID-19: a case-control study. *Odontology*. 2023. p. 1–13.
- Costa CA, Vilela ACS, Oliveira SA, Gomes TD, Andrade AAC, Leles CR, Costa NL. Poor oral health status and adverse COVID-19 outcomes: a preliminary study in hospitalized patients. *J Periodontol*. 2022;93(12):1889–901.
- Guardado-Luevanos I, Bologna-Molina R, Zepeda-Nuño JS, Isordia-Espinoza M, Molina-Frechero N, González-González R, Pérez-Pérez M, López-Verdín S. Self-Reported Periodontal Disease and Its Association with SARS-CoV-2 Infection. *Int J Environ Res Public Health*. 2022;19(16):10306.
- Gujar D, Joshi C, Pandya DJ, Nayak K, Shafiuddin M, Sameerudeen S, Mahajan A. Exploring the correlation between covid-19 and periodontal diseases-An original research. *Eur J Mol Clin Med*. 2022;9(7):8707–12.
- Holt H, Talaei M, Greenig M, Zenner D, Symons J, Relton C, et al. Risk factors for developing COVID-19: a population-based longitudinal study (COVIDENCE UK). *Thorax*. 2022;77(9):900–12.
- Kalsi R, Ahmad Z, Siddharth M, Vandana K, Arora S, Saurav K. Correlation of COVID-19 with severity of periodontitis-A clinical and biochemical study. *Indian J Dent Res*. 2022;33(3):307–12.

41. Kamel AHM, Basuoni A, Salem ZA, AbuBakr N. The impact of oral health status on COVID-19 severity, recovery period and C-reactive protein values. *Br Dent J*. 2021. p. 1–7.
42. Kaur A, Sandhu HS, Sarwal A, Bhagat S, Dodwad R, Singh G, Gambhir RS. Assessment of correlation of COVID-19 infection and periodontitis-A comparative study. *J Family Med Prim Care*. 2022;11(5):1913–7.
43. Koppolu P, Genady EM, Albdeirat LM, Sebai FA, Alrashdi DM, Lingam AS, FA RA, Al-Khalifa FI, Abdelrahim RK: Association between severity of COVID-19, Periodontal health and disease in Riyadh subpopulation. *Int J Mycobacteriol*. 2023;12(1):33–7.
44. Mishra S, Gupta V, Rahman W, Gazala M, Anil S. Association between Periodontitis and COVID-19 Based on Severity Scores of HRCT Chest Scans. *Dent J*. 2022;10(6):106.
45. Poyato-Borrego M, León-López M, Martín-González J, Cisneros-Herreros JM, Cabanillas-Balsera D, Segura-Egea JJ. Endodontic variables in patients with SARS-CoV-2 infection (COVID-19) in relation to the severity of the disease. *Med Oral Patol Oral Cir Bucal*. 2023;28(4):e355–e361.
46. Sari A, Dikmen NK, Nibali L. Association between periodontal diseases and COVID-19 infection: a case-control study with a longitudinal arm. *Odontology*. 2023. p. 1–9.
47. Sirin DA, Ozcelik F. The relationship between COVID-19 and the dental damage stage determined by radiological examination. *Oral Radiol*. 2021;37(4):600–9.
48. Takahashi Y, Watanabe N, Kamio N, Kobayashi R, Iinuma T, Imai K. Aspiration of periodontopathic bacteria due to poor oral hygiene potentially contributes to the aggravation of COVID-19. *J Oral Sci*. 2020;63(1):1–3.
49. MadapusiBalaji T, Varadarajan S, Rao USV, Raj AT, Patil S, Arakeri G, Brennan PA. Oral cancer and periodontal disease increase the risk of COVID 19? A mechanism mediated through furin and cathepsin overexpression. *Med Hypotheses*. 2020;144: 109936.
50. Chakraborty S. Metagenome of SARS-Cov2 patients in Shenzhen with travel to Wuhan shows a wide range of species-Lautropia, Cutibacterium, Haemophilus being most abundant-and Campylobacter explaining diarrhea. Available at: <https://osf.io/jegwq>, <https://doi.org/10.31219/osf.io/jegwq>. Accessed June 2023.
51. Gomes-Filho IS, Cruz SSD, Trindade SC, Passos-Soares JS, Carvalho-Filho PC, Figueiredo A, Lyrio AO, Hintz AM, Pereira MG, Scannapieco F. Periodontitis and respiratory diseases: a systematic review with meta-analysis. *Oral Dis*. 2020;26(2):439–46.
52. Kim SJ, Kim K, Choi S, Chang J, Kim SM, Park SM, Cho HJ. Chronic periodontitis and community-acquired pneumonia: a population-based cohort study. *BMC Pulm Med*. 2019;19(1):268.
53. Hata R, Noguchi S, Kawanami T, Yamasaki K, Akata K, Ikegami H, Fukuda K, Hirashima S, Miyawaki A, Fujino Y, et al. Poor oral hygiene is associated with the detection of obligate anaerobes in pneumonia. *J Periodontol*. 2020;91(1):65–73.
54. Sakaguchi W, Kubota N, Shimizu T, Saruta J, Fuchida S, Kawata A, Yamamoto Y, Sugimoto M, Yakeishi M, Tsukinoki K. Existence of SARS-CoV-2 Entry Molecules in the Oral Cavity. *Int J Mol Sci*. 2020;21(17):6000.
55. Xu H, Zhong L, Deng J, Peng J, Dan H, Zeng X, Li T, Chen Q. High expression of ACE2 receptor of 2019-nCoV on the epithelial cells of oral mucosa. *Int J Oral Sci*. 2020;12(1):8.
56. Natto ZS, Afeef M, Bakhrebah MA, Ashi H, Alzahrani KA, Alhethel AF, Fletcher HM. Can periodontal pockets and caries lesions act as reservoirs for coronavirus? *Mol Oral Microbiol*. 2022;37(2):77–80.
57. Sekundo C, Stock C, Jürges H, Listl S. Patients' self-reported measures of oral health-A validation study on basis of oral health questions used in a large multi-country survey for populations aged 50. *Gerodontology*. 2019;36(2):171–9.
58. Metelli S, Chaimani A. Challenges in meta-analyses with observational studies. *Evid Based Ment Health*. 2020;23(2):83–7.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

